

Supplementary appendix

Supplement to: Ho ISS, Azcoaga-Lorenzo A, Akbari A, et al. Variation in the estimated prevalence of multimorbidity: systematic review and meta-analysis of 193 studies.

Table of Contents

Table S1: Search strategy.....	3
Table S2: Summary of the characteristics of outlying studies.....	4
Table S3: Characteristics of 24 outlying studies.....	5
Table S4: Characteristics of 193 included studies	9
Table S5: Associations between predictors	25
Table S6: Risk of bias assessment of included studies	26
Table S7: Output of adjusted meta-analytic model based on 217 studies	39
Table S8: Definition of variables	40
Figure S1: Graphical display of study effect sizes and heterogeneity	41
Figure S2: Process of examining and identifying outlying studies in meta-analysis	42
Figure S3: Summary of risk of bias assessment	43
Figure S4: Meta-regression trees for predicting the pooled estimated prevalence of multimorbidity (based on ‘mean age’ and ‘number of conditions’ predictors. unit: log(odds)).....	44
References.....	45

Table S1: Search strategy

Database	Search strategy
Ovid Interface PsycINFO Embase Global Health Ovid MEDLINE	<ol style="list-style-type: none"> 1. (multimorbidit\$ or multi-morbidit\$ or comorbidit\$ or co-morbidit\$ or polymorbidit\$ or poly-morbidit\$ or multicondition\$ or multicondition\$ or "multiple chronic condition\$" or "morbidity burden" or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) adj2 (disease\$ or illness\$ or condition\$ or diagnos\$ or morbid\$))).m_titl. 2. (measure\$ or index or indices or instrument\$ or scale\$ or "disease count\$").mp. 3. 1 and 2 4. Limit 3 to human
EBSCO Interface CINAHL Plus	<ol style="list-style-type: none"> 1. MM (multimorbidit* or multi-morbidit* or comorbidit* or co-morbidit* or polymorbidit* or poly-morbidit* or multicondition* or multicondition* or "multiple chronic condition*" or "morbidity burden" or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) N2 (disease* or illness* or condition* or diagnos* or morbid*))) 2. AB (measure* or index or indices or instrument* or scale*) 3. 1 AND 2 Limiters – Full Text; Human; Language: English
Scopus	TITLE (multimorbidit* or multi-morbidit* or comorbidit* or co-morbidit* or polymorbidit* or poly-morbidit* or multicondition* or multicondition* or "multiple chronic condition*" or "morbidity burden" or ((multiple or coexisting or co-existing or concurrent or con-current or morbid or co-morbid) W/2 (disease* or illness* or condition* or diagnos?s or morbid*))) AND TITLE (measure* or index or indices or instrument* or scale* or "disease counts")
Web of Science	(TI=(measure* or index or indices or instrument* or scale*))AND (TI=(multimorbidit* or multi-morbidit* or comorbidit* or co-morbidit* or polymorbidit* or poly-morbidit* or multicondition* or multicondition* or 'multiple chronic condition*' or 'morbidity burden' or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) NEAR/2 (disease* or illness* or condition* or diagnos* or morbid*)))) AND LANGUAGE: (English)
Cochrane library	(multimorbidity or multi-morbidity or comorbidity or co-morbidity or polymorbidity or poly-morbidity or multicondition or multicondition or 'multiple chronic conditions' or 'morbidity burden' or ((multiple or coexisting or co-existing or concurrent or con-current or comorbid or co-morbid) NEAR/2 (disease or illness or condition or diagnosis or morbid))) AND (measure or index or indices or instrument or scale or "disease count*"):ti
ProQuest Dissertations & Theses Global	ti((multimorbidit* OR multi-morbidit* OR comorbidit* OR co-morbidit* OR polymorbidit* OR poly-morbidit* OR multicondition* OR multicondition* OR 'multiple chronic condition*' OR 'morbidity burden' OR ((multiple OR coexisting OR co-existing OR concurrent OR con-current OR morbid OR co-morbid) NEAR/2 (disease* OR illness* OR condition* OR diagnos?s OR morbid*)))) AND noft((measure* OR index OR indices OR instrument* OR scale*)) Limited by: Manuscript type: Doctoral dissertations, Master's theses Language: English

Table S2: Summary of the characteristics of outlying studies

Name of variable	Outlying studies (n=24)	All studies (n=217)
Prevalence of multimorbidity (%)	Range: 7.3 to 89.1 Pooled prevalence with the REML estimator: 31.0 (21.6–42.2)	Range: 2.7–95.6 Pooled prevalence with the REML estimator: 41.1 (37.7–44.6)
Mean age of study population (year)	Range of mean age: 39.6 to 82.2 Median of mean age: 56.6 (Q1, Q3: 52.3, 66.4)	Range of mean age: 32.2 to 83.8 Median of mean age: 62.4 (Q1, Q3: 50.2, 72.0)
No of conditions (count)	Range: 7 to 259 Median: 34 (Q1, Q3: 19.5, 54.5)	Range: 3 to 259 Median: 14.0 (Q1, Q3: 9, 21)
Country income (count, %)		
High income	21 (87.5%)	166 (76.5%)
Low- or Middle-income	3 (11.5%)	51 (23.5%)
Continent (count, %)		
Europe	6 (25.0%)	70 (32.3%)
North America	7 (29.2%)	54 (24.9%)
Asia	7 (29.2%)	51 (23.5%)
Australasia	3 (12.5%)	14 (6.5%)
Multiple continents	1 (4.2%)	12 (5.5%)
South America		12 (5.5%)
Africa		4 (1.8%)
Study population (count, %)		
Only older people	2 (8.3%)	65 (30.0%)
Middle-aged and older	1 (4.2%)	47 (21.7%)
All adults	15 (62.5%)	99 (45.6%)
Only children	1 (4.2%)	1 (0.5%)
All age population	5 (20.8%)	5 (2.3%)
Setting (count, %)		
Community	12 (50.0%)	159 (73.3%)
Primary care	7 (29.2%)	39 (18.0%)
Hospital	4 (16.7%)	18 (8.3%)
Care home	1 (4.2%)	1 (0.5%)
Source (count, %)		
Self-report	8 (33.3%)	158 (72.8%)
Database	16 (66.6%)	59 (27.2%)
Risk of bias assessment (count, %)		
Low	4 (16.7%)	13 (6.0%)
Moderate	19 (79.2%)	181 (83.4%)
High	1 (4.2%)	23 (10.6%)

IQR: Interquartile range. SD: Standard deviation. The percentages were rounded so they do not add up to 100%.

Table S3: Characteristics of 24 outlying studies

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias	Rationale for exclusion
¹ Stanley et al (2018)	New Zealand	Australasia	High	Hospitals	All adults	Not reported	3489747	Medical records and administrative database	61	275706	0.08	Moderate	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) and the studentized residual of this study is more than 2 standard deviations away from its expected value.
² Lenzi et al (2016)	Italy	Europe	High	Hospitals	All adults	66.4	3759836	Medical records and administrative database	26	574208	0.15	Moderate	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
³ Hu et al (2019)	Taiwan	Asia	High	Community	All adults	Not reported	1429527	Medical records and administrative database	20	939485	0.66	Low	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
⁴ Gawron et al (2020)	USA	North America	High	Hospitals	All adults but not older people	Not reported	741612	Medical records and administrative database	Not reported	53824	0.07	Moderate	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis) and the studentized residual of this study is more than 2 standard deviations away from its expected value.
⁵ Low et al (2019)	Singapore	Asia	High	Community	All adults	39.6	1181024	Self-report	48	309428	0.26	Low	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
⁶ Wang et al (2014)	China	Asia	Low or middle	Community	Whole population	Not reported	162464	Self-report	40	17987	0.11	Low	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
⁷ Gaulin et al (2019)	Canada	North America	High	Hospitals	All adults	51.2	1316832	Medical records and administrative database	34	416282	0.32	Moderate	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias	Rationale for exclusion
⁸ Violan et al (2014)	Spain	Europe	High	Primary care	All adults	47.4	1356761	Medical records and administrative database	146	645818	0.48	Moderate	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
⁹ Nicholson et al (2019)	Canada	North America	High	Primary care	All adults	52.3	367743	Medical records and administrative database	20	195838	0.53	High	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
¹⁰ Bao et al (2019)	China	Asia	Low or middle	Community	Middle aged and older	61.36	18137	Self-report	19	3773	0.21	Moderate	Contributing to high levels of heterogeneity of effect sizes (Leave-one-out analysis)
¹¹ Fortin et al (2005)	Canada	North America	High	Primary care	All adults	56.55	980	Medical records and administrative database	14	873	0.89	Moderate	The studentized residual of this study is more than 2 standard deviations away from its expected value.
¹² Prazeres et al (2015)	Portugal	Europe	High	Primary care	All adults	56.3	1993	Medical records and administrative database	147	1449	0.73	Moderate	Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection)
¹³ Lawson et al (2013)	UK	Europe	High	Community	All adults	72.7	7054	Medical records and administrative database	40	1243	0.18	Moderate	Irregular patterns found in compositional data (in scatter plot and Mahalanobis distance test)- low prevalence in studies with high mean participant age and a larger number of conditions
¹⁴ Sullivan et al (2012)	USA	North America	High	Community	All adults	Not reported	47178	Medical records and administrative database	259	19666	0.42	Moderate	Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection)

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias	Rationale for exclusion
¹⁵ Peng et al (2020)	China	Asia	Low or middle	Community	Only older people	71.6	1321	Self-report	15	589	0.45	Moderate	Contributing to high levels of heterogeneity of effect sizes (in leave-one-out analysis)
¹⁶ Excoffier et al (2018)	Switzerland	Europe	High	Primary care	All adults	56.5	2904	Medical records and administrative database	75	1513	0.52	Moderate	Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection)
¹⁷ Chung et al (2015)	Hong Kong	Asia	High	Community	All adults	Not reported	25780	Self-report	46	3227	0.13	Moderate	Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection)
¹⁸ Ki et al (2017)	South Korea	Asia	High	Community	All adults	57.05	19942	Medical records and administrative database	66	5979	0.30	Moderate	Its Mahalanobis distance exceeds the chi-squared critical value at a 0.01 significance level (multivariate outlier detection)
¹⁹ Bobo et al (2016)	USA	North America	High	Community	Whole population	Not reported	138858	Self-report	19	33682	0.24	Moderate	Infrequent values in compositional categorical data (few studies focused on whole population)
²⁰ Randall et al (2018)	Australia	Australasia	High	Community	Whole population	Not reported	5437018	Self-report	30	660449	0.12	Moderate	Infrequent values in compositional categorical data (few studies focused on whole population)
²¹ Russell et al (2020)	New Zealand	Australasia	High	Community	Only children	Not reported	3838	Self-report	7	374	0.10	Moderate	Infrequent values in compositional categorical data (only one study focused on children population)
²² Barnett et al (2012)	UK	Europe	High	Primary care	Whole population	Not reported	1751841	Medical records and administrative database	40	406427	0.23	Low	Infrequent values in compositional categorical data (few studies focused on whole population)

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias	Rationale for exclusion
²³ St Sauver et al (2015)	USA	North America	High	Primary care	Whole population	Not reported	106061	Medical records and administrative database	20	34592	0.33	Moderate	Infrequent values in compositional categorical data (few studies focused on whole population)
²⁴ Vetrano et al (2016)	Denmark, Finland, Iceland, Italy, the Netherlands, Norway, United Kingdom, Czech Republic, France, Sweden and Germany, Canada	Multiple continents	High	Care homes	Only older people	82.2	6903	Medical records and administrative database	13	5098	0.74	Moderate	Infrequent values in compositional categorical data (only one study focused on care home)

MM: Multimorbidity. No of participants: The total number of participants in the denominator for estimating prevalence in a study (which could be a subset in some included studies)

Table S4: Characteristics of 193 included studies

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
²⁵ Aarts et al (2012)	The Netherlands	Europe	High	Primary care	All adults	55.4	1184	Medical records and administrative database	23	420	0.35	Moderate
²⁶ Aarts et al (2011a)	The Netherlands	Europe	High	Community	Middle aged and older	70	15188	Self-report	Not reported	7729	0.51	Moderate
²⁷ Aarts et al (2011b)	The Netherlands	Europe	High	Primary care	All adults	55.4	1763	Medical records and administrative database	23	985	0.56	Moderate
²⁸ Abizanda et al (2014)	Spain	Europe	High	Primary care	Only older people	78.6	842	Medical records and administrative database	14	580	0.69	Moderate
²⁹ Agborsangaya et al (2012)	Canada	North America	High	Community	All adults	46.6	4003	Self-report	16	919	0.23	Moderate
³⁰ Agborsangaya et al (2013)	Canada	North America	High	Community	All adults	47.8	4803	Self-report	16	1729	0.36	Moderate
³¹ Agborsangaya et al (2014)	Canada	North America	High	Community	All adults	47.7	4752	Self-report	16	1597	0.34	Moderate
³² Ahrenfeldt et al (2019)	Europe	Europe	High	Community	Middle aged and older	66.25	244258	Self-report	10	90652	0.37	Moderate
³³ Alimohammadian et al (2017)	Iran	Asia	Low or middle	Community	Middle aged and older	Not reported	49946	Self-report	8	10035	0.20	Moderate
³⁴ Angst et al (2002)	Switzerland	Europe	High	Primary care	All adults	Not reported	591	Medical records and administrative database	10	201	0.34	High

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
³⁵ Appa et al (2014)	USA	North America	High	Community	All adults	60.2	1997	Self-report	16	1417	0.71	Moderate
³⁶ Adams et al (2017)	USA	North America	High	Community	All adults	Not reported	400000	Self-report	12	191600	0.48	Moderate
³⁷ Ahmadi et al (2016)	Iran	Asia	Low or middle	Community	Middle aged and older	52.1	49946	Self-report	8	10035	0.20	Moderate
³⁸ Amaral et al (2018)	Brazil	South America	Low or middle	Community	Only older people	Not reported	264	Self-report	8	175	0.66	Moderate
³⁹ An et al (2016)	South Korea	Asia	High	Community	Middle aged and older	54.8	10118	Self-report	8	3228	0.32	Moderate
⁴⁰ Araujo et al (2018)	Brazil	South America	Low or middle	Community	All adults	Not reported	4001	Self-report	12	1160	0.29	Moderate
⁴¹ Arnold-Reed et al (2018)	Australia	Australasia	High	Primary care	All adults	38.2	4285	Medical records and administrative database	43	2269	0.53	Moderate
⁴² Arokiasamy et al (2015)	6 low middle income countries (China, Ghana, India, Mexico, Russia, South Africa)	Multiple continents	Low or middle	Community	All adults	Not reported	42236	Self-report	8	9250	0.22	Moderate
⁴³ Sinnige et al (2015)	The Netherlands	Europe	High	Primary care	Middle aged and older	66.9	120480	Medical records and administrative database	29	74733	0.62	Moderate
⁴⁴ Zemedikun et al (2018)	UK	Europe	High	Community	Middle aged and older	Median age 58	502643	Medical records and administrative database	36	95710	0.19	Moderate
⁴⁵ Wensing et al (2001)	The Netherlands	Europe	High	Primary care	All adults	Not reported	3867	Self-report	25	626	0.16	Moderate
⁴⁶ Mounce et al (2018)	UK	Europe	High	Community	Middle aged and older	Not reported	4564	Self-report	15	1553	0.34	Moderate
⁴⁷ Taylor et al (2010)	Australia	Australasia	High	Community	All adults	Not reported	3206	Self-report	7	547	0.17	Low

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
⁴⁸ Vancampfort et al (2019)	Six low and middle income countries (China, Ghana, India, Mexico, Russia, and South Africa)	Multiple continents	Low or middle	Community	Middle aged and older	62.4	34129	Self-report	11	15529	0.46	Moderate
⁴⁹ Vancampfort et al (2018)	Six low and middle income countries (China, Ghana, India, Mexico, Russia, and South Africa)	Multiple continents	Low or middle	Community	Only older people	72.6	14585	Self-report	11	8780	0.60	Moderate
⁵⁰ Aubert et al (2016)	Switzerland	Europe	High	Primary care	Middle aged and older	63.5	1002	Medical records and administrative database	17	676	0.67	Moderate
⁵¹ Autenrieth et al (2013)	Germany	Europe	High	Community	Only older people	75.7	1007	Self-report	13	658	0.65	Moderate
⁵² Bahler et al (2015)	Switzerland	Europe	High	Community	Only older people	74.9	229493	Medical records and administrative database	22	175752	0.77	Moderate
⁵³ Vancampfort et al (2017)	44 low and middle income countries	Multiple continents	Low or middle	Community	All adults	38.3	194431	Self-report	11	27518	0.14	Moderate
⁵⁴ Banjare et al (2014)	India	Asia	Low or middle	Community	Only older people	Not reported	310	Self-report	20	176	0.57	Moderate
⁵⁵ Barra et al (2015)	USA	North America	High	Community	All adults	45.36	43079	Self-report	Not reported	22412	0.52	Moderate
⁵⁶ Bernard et al (2016)	Australia	Australasia	High	Hospitals	Only older people	81.8	306	Medical records and administrative database	19	125	0.41	High
⁵⁷ Biswas et al (2019)	Bangladesh	Asia	Low or middle	Community	All adults	Not reported	8763	Self-report	3	1078	0.12	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
⁵⁸ Blakemore et al (2016)	UK	Europe	High	Primary care	Only older people	75	4377	Self-report	24	2631	0.60	Moderate
⁵⁹ Blyth et al (2008)	Australia	Australasia	High	Community	Only older people	76.9	1685	Self-report	18	920	0.55	Moderate
⁶⁰ Bowling et al (2019)	USA	North America	High	Community	Middle aged and older	56.7	4217	Self-report	12	3053	0.72	Moderate
⁶¹ Britt et al (2008)	Australia	Australasia	High	Primary care	All adults	Not reported	9156	Medical records and administrative database	18	3398	0.37	Moderate
⁶² Broeiro-Goncalves et al (2019)	Portugal	Europe	High	Hospitals	All adults	59.8	800376	Medical records and administrative database	22	335357	0.42	Moderate
⁶³ Bruce et al (2010)	Canada	North America	High	Community	All adults	37.8	453	Self-report	4	163	0.36	High
⁶⁴ Burgers et al (2010)	France, Germany, Canada, Australia, Netherlands, New Zealand, UK, USA	Multiple continents	High	Community	All adults	Not reported	8973	Self-report	7	4037	0.45	Moderate
⁶⁵ Burke et al (2017)	US, Europe, Asia	Multiple continents	High	Community	Only older people	Not reported	4668	Self-report	9	2165	0.46	Moderate
⁶⁶ Buurman et al (2016)	The Netherlands	Europe	High	Hospitals	Only older people	78.2	639	Medical records and administrative database	35	440	0.69	Moderate
⁶⁷ Calderon-Larranaga et al (2017)	Sweden	Europe	High	Primary care	Only older people	74.6	3363	Self-report	60	2980	0.89	Moderate
⁶⁸ Camargo-Casas et al (2018)	Colombia	South America	Low or middle	Community	Only older people	71.1	2000	Self-report	12	808	0.40	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
⁶⁹ Canevelli et al (2019)	Italy	Europe	High	Primary care	Only older people	75.1	185	Medical records and administrative database	18	162	0.88	High
⁷⁰ Chamberlain et al (2020)	USA	North America	High	Community	All adults	Not reported	198941	Self-report	21	78527	0.39	Low
⁷¹ Chen et al (2018)	China	Asia	Low or middle	Community	Only older people	Not reported	30774	Medical records and administrative database	33	25101	0.82	Low
⁷² Chen et al (2018)	China	Asia	Low or middle	Community	Middle aged and older	Not reported	3737	Self-report	16	1722	0.46	Moderate
⁷³ Cheung et al (2013)	Hong Kong (SAR of China)	Asia	High	Community	Middle aged and older	71.3	1145	Self-report	18	654	0.57	Moderate
⁷⁴ Chu et al (2018)	Hong Kong (SAR of China)	Asia	High	Primary care	Middle aged and older	Not reported	382	Medical records and administrative database	40	206	0.54	Moderate
⁷⁵ Chudasama et al (2019)	UK	Europe	High	Community	Middle aged and older	Median age:58	491939	Medical records and administrative database	36	96622	0.20	Moderate
⁷⁶ Cimarras-Otal et al (2014)	Spain	Europe	High	Community	All adults	Not reported	22190	Self-report	20	7830	0.35	Moderate
⁷⁷ Chin et al (2016)	Hong Kong (SAR of China)	Asia	High	Primary care	All adults	Median age: 48	9259	Self-report	8	2350	0.25	Moderate
⁷⁸ Agrawal et al (2016)	India, China, Russia, Mexico, South Africa, Ghana	Multiple continents	Low or middle	Community	All adults	57.8	40166	Self-report	9	9238	0.23	Moderate
⁷⁹ Gu et al (2018)	China	Asia	Low or middle	Community	Only older people	Not reported	411	Self-report	13	232	0.56	Moderate
⁸⁰ Gunn et al (2012)	Australia	Australasia	High	Primary care	All adults	50.89	6864	Self-report	12	2154	0.31	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
⁸¹ Han et al (2013)	USA	North America	High	Primary care	Only older people	76	159	Medical records and administrative database	18	117	0.74	High
⁸² Hanlon et al (2018)	UK	Europe	High	Community	All adults	Not reported	493737	Medical records and administrative database	42	161576	0.33	Low
⁸³ Jantsch et al (2018)	Brazil	South America	Low or middle	Community	All adults	42	3092	Self-report	11	912	0.29	Moderate
⁸⁴ John et al (2003)	USA	North America	High	Community	Only older people	71.3	992	Self-report	11	732	0.74	High
⁸⁵ Johnson-Lawrence et al (2017)	USA	North America	High	Community	All adults	49.9	115097	Self-report	9	27278	0.24	Moderate
⁸⁶ Johnston et al (2019)	UK	Europe	High	Community	All adults	48	7184	Self-report	Not reported	388	0.05	Moderate
⁸⁷ Jones et al (2016)	USA	North America	High	Community	Only older people	Not reported	6964	Self-report	10	4951	0.71	Moderate
⁸⁸ Jovic et al (2016)	Serbia	Europe	Low or middle	Community	All adults	49.4	13103	Self-report	13	3522	0.27	Moderate
⁸⁹ Juul-Larsen et al (2020)	Denmark	Europe	High	Hospitals	Only older people	Median age: 78	369	Self-report	34	311	0.84	Moderate
⁹⁰ Hudon et al (2008)	Canada	North America	High	Community	All adults	Not reported	16782	Self-report	25	5343	0.32	Low
⁹¹ Hussain et al (2015)	Indonesia	Asia	Low or middle	Community	Middle aged and older	Not reported	9438	Self-report	12	3369	0.36	Moderate
⁹² Ie et al (2017)	USA	North America	High	Hospitals	Only older people	Not reported	1084	Medical records and administrative database	24	1036	0.96	High
⁹³ Ishizaki et al (2019)	Japan	Asia	High	Community	Only older people	76.9	2525	Self-report	9	1121	0.44	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
⁹⁴ Danon-Hersch et al (2012)	Switzerland	Europe	High	Community	Only older people	Not reported	1283	Self-report	12	448	0.35	Moderate
⁹⁵ de Heer et al (2013)	USA	North America	High	Community	All adults	47.72	1002	Self-report	19	378	0.38	Moderate
⁹⁶ Demirchyan et al (2013)	Armenia	Asia	Low or middle	Community	All adults	58.8	721	Self-report	Not reported	564	0.78	High
⁹⁷ Fabbri et al (2015)	Italy	Europe	High	Community	Only older people	73.6	1018	Self-report	15	458	0.45	Moderate
⁹⁸ Fillenbaum et al (2000)	USA	North America	High	Community	Only older people	73.44	4034	Self-report	5	1181	0.29	Moderate
⁹⁹ Kaneko et al (2019)	Japan	Asia	High	Community	Only older people	Not reported	253	Self-report	Not reported	135	0.53	Moderate
¹⁰⁰ Kang et al (2017)	South Korea	Asia	High	Primary care	All adults	32.2	590	Medical records and administrative database	14	153	0.26	Moderate
¹⁰¹ Gandhi et al (2020)	USA	North America	High	Community	All adults	Not reported	9499	Self-report	8	3379	0.36	Moderate
¹⁰² Costa et al (2018)	Brazil	South America	Low or middle	Community	Only older people	Not reported	1451	Self-report	29	1343	0.93	Moderate
¹⁰³ Rizzuto et al (2017)	Sweden	Europe	High	Community	Only older people	Not reported	1099	Self-report	36	774	0.70	Moderate
¹⁰⁴ Dhalwani et al (2017)	UK	Europe	High	Community	Middle aged and older	Not reported	5476	Self-report	18	1156	0.21	Moderate
¹⁰⁵ Elixhauser et al (1998)	USA	North America	High	Hospitals	All adults	57.1	1779167	Medical records and administrative database	30	619150	0.35	Low
¹⁰⁶ Fabbri et al (2015)	USA	North America	High	Hospitals	Only older people	72.3	695	Self-report	15	440	0.63	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹⁰⁷ Fortin et al (2014)	Canada	North America	High	Community	Middle aged and older	57.8	1196	Self-report	14	599	0.50	Moderate
¹⁰⁸ Fuchs et al (1998)	Israel	Asia	High	Community	Only older people	Not reported	1820	Self-report	14	1174	0.65	Moderate
¹⁰⁹ Galenkamp et al (2011)	The Netherlands	Europe	High	Community	Middle aged and older	69.2	2046	Self-report	7	876	0.43	High
¹¹⁰ Galenkamp et al (2016)	Germany, UK, Italy, The Netherlands, Spain and Sweden	Europe	High	Community	Only older people	74.2	2792	Self-report	8	1358	0.49	Moderate
¹¹¹ Gamma et al (2001)	Switzerland	Europe	High	Community	All adults	Not reported	407	Self-report	14	53	0.13	High
¹¹² Ge et al (2018)	Singapore	Asia	High	Community	All adults	51.4	1940	Self-report	17	715	0.37	Moderate
¹¹³ Ge et al (2019)	Singapore	Asia	High	Community	All adults	51.3	1932	Self-report	17	564	0.29	Moderate
¹¹⁴ Gould et al (2016)	USA	North America	High	Community	Only older people	74.82	4184	Self-report	7	2932	0.70	Moderate
¹¹⁵ Habib et al (2014)	Lebanon	Asia	Low or middle	Community	All adults	46.6	2501	Self-report	Not reported	665	0.27	Moderate
¹¹⁶ Harrison et al (2017)	Australia	Australasia	High	Primary care	All adults	Not reported	8707	Medical records and administrative database	28	2838	0.33	Moderate
¹¹⁷ Hayek et al (2017)	Israel	Asia	High	Community	All adults	47.2	4325	Self-report	10	1579	0.37	Moderate
¹¹⁸ Henninger et al (2012)	USA	North America	High	Community	Only older people	76	3212	Self-report	9	1753	0.55	Moderate
¹¹⁹ Hernandez et al (2019)	Ireland	Europe	High	Community	Middle aged and older	Not reported	6101	Self-report	31	4468	0.73	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹²⁰ Ho et al (2014)	Singapore	Asia	High	Community	Middle aged and older	66.15	1844	Self-report	12	830	0.45	Moderate
¹²¹ Khan et al (2019)	Bangladesh	Asia	Low or middle	Community	All adults	58.6	12338	Self-report	6	1031	0.08	Low
¹²² Kiliari et al (2013)	Cyprus	Europe	High	Community	All adults	53	465	Self-report	Not reported	132	0.28	Moderate
¹²³ King et al (2018)	USA	North America	High	Community	All adults	Not reported	5541	Self-report	11	3342	0.60	Moderate
¹²⁴ Kingston et al (2018)	UK	Europe	High	Community	All adults	Not reported	9723900	Self-report	12	5250906	0.54	High
¹²⁵ Koyanagi et al (2018)	China, Ghana, India, Mexico, Russia, and South Africa	Multiple continents	Low or middle	Community	Middle aged and older	62.1	32715	Self-report	10	16324	0.50	Moderate
¹²⁶ Kriegsman et al (2004)	The Netherlands	Europe	High	Community	Middle aged and older	69.2	2489	Self-report	7	519	0.21	Moderate
¹²⁷ Kristensen et al (2019)	Germany	Europe	High	Community	Middle aged and older	63.47	19605	Self-report	13	12600	0.64	Moderate
¹²⁸ Kristensen et al (2019)	Germany	Europe	High	Community	Middle aged and older	64.37	7604	Self-report	13	5140	0.68	Moderate
¹²⁹ Kunna et al (2017)	China, Ghana	Multiple continents	Low or middle	Community	Middle aged and older	Not reported	15864	Self-report	7	4731	0.30	Low
¹³⁰ Kuwornu et al (2014)	Canada	North America	High	Community	All adults	51.05	3284	Self-report	15	1143	0.35	Moderate
¹³¹ Lai et al (2019)	Hong Kong (SAR of China)	Asia	High	Community	All adults	Not reported	69636	Self-report	14	3898	0.06	Moderate
¹³² Lai et al (2018)	Hong Kong (SAR of China)	Asia	High	Community	All adults	Not reported	300	Self-report	11	48	0.16	Moderate
¹³³ Laires et al (2019)	Portugal	Europe	High	Community	All adults	Not reported	15196	Self-report	13	6671	0.44	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹³⁴ Lang et al (2015)	USA	North America	High	Community	Middle aged and older	53.4	3058	Self-report	6	948	0.31	Moderate
¹³⁵ Le Cossec et al (2016)	France	Europe	High	Community	Middle aged and older	70	15325	Self-report	4	3528	0.23	Moderate
¹³⁶ Lee et al (2007)	USA	North America	High	Hospitals	Middle aged and older	Not reported	741847	Medical records and administrative database	11	302792	0.41	Low
¹³⁷ Lee et al (2018)	Taiwan	Asia	High	Community	Only older people	Not reported	20898	Medical records and administrative database	Not reported	4234	0.20	High
¹³⁸ Li et al (2016)	UK	Europe	High	Primary care	All adults	Not reported	27806	Self-report	12	10332	0.37	Moderate
¹³⁹ Li et al (2019)	USA	North America	High	Community	Middle aged and older	67.4	14996	Self-report	8	9805	0.65	Moderate
¹⁴⁰ Lujic et al (2017)	Australia	Australasia	High	Community	Middle aged and older	70.2	90352	Self-report	8	33792	0.37	Moderate
¹⁴¹ Lupianez-Villanueva et al (2018)	14 European countries	Europe	High	Community	All adults	Not reported	14000	Self-report	13	3416	0.24	Moderate
¹⁴² Zhou et al (2018)	Bangladesh, India and China	Asia	Low or middle	Community	All adults	Not reported	18696	Self-report	9	3512	0.19	Moderate
¹⁴³ Zhang et al (2019)	China	Asia	Low or middle	Community	Only older people	70.5	11707	Self-report	11	5104	0.44	Moderate
¹⁴⁴ Wong et al (2010)	Canada	North America	High	Community	Only older people	Not reported	740	Self-report	7	489	0.66	Moderate
¹⁴⁵ Weimann et al (2016)	South Africa	Africa	Low or middle	Community	All adults	34	18526	Self-report	4	506	0.027	Moderate
¹⁴⁶ Wang et al (2017)	Australia	Australasia	High	Community	All adults	44	8820	Self-report	8	2539	0.29	Moderate
¹⁴⁷ Wang et al (2019)	South Africa	Africa	Low or middle	Community	Only older people	Not reported	2627	Self-report	5	439	0.17	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹⁴⁸ Wade et al (2019)	New Zealand	Australasia	High	Community	All adults	59.05	7654	Self-report	12	2786	0.36	Moderate
¹⁴⁹ Maciejewski et al (2019)	USA	North America	High	Community	Only older people	77.1	20124230	Medical records and administrative database	19	14425446	0.72	Moderate
¹⁵⁰ Marengoni et al (2016)	Sweden	Europe	High	Community	Only older people	74.4	3155	Medical records and administrative database	14	1654	0.52	Moderate
¹⁵¹ Marengoni et al (2009)	Sweden	Europe	High	Community	Only older people	Not reported	1099	Self-report	22	575	0.52	Moderate
¹⁵² Marques et al (2018)	13 European countries	Europe	High	Community	All adults	50.2	32931	Self-report	6	7113	0.22	Moderate
¹⁵³ Mavaddat et al (2014)	UK	Europe	High	Primary care	Middle aged and older	58.7	11439	Self-report	6	1006	0.09	Moderate
¹⁵⁴ McDaid et al (2013)	Ireland	Europe	High	Community	Middle aged and older	Not reported	6018	Self-report	8	733	0.12	High
¹⁵⁵ Melis et al (2014)	Sweden	Europe	High	Hospitals	Only older people	83.75	390	Medical records and administrative database	39	213	0.55	Moderate
¹⁵⁶ Min et al (2007)	USA	North America	High	Community	Only older people	81	372	Self-report	9	230	0.62	High
¹⁵⁷ Momtaz et al (2010)	Malaysia	Asia	High	Community	Only older people	69.26	385	Self-report	16	165	0.43	Moderate
¹⁵⁸ Mondor et al (2018)	Canada	North America	High	Community	All adults	Not reported	27195	Medical records and administrative database	17	11390	0.42	Moderate
¹⁵⁹ Muggah et al (2012)	Canada	North America	High	Community	All adults	Not reported	28450000	Medical records and administrative database	9	4523550	0.16	Moderate
¹⁶⁰ Nagel et al (2008)	Germany	Europe	High	Community	Middle aged and older	56.5	13781	Self-report	15	9275	0.67	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹⁶¹ Niedzwiedz et al (2019)	USA	North America	High	Community	Middle aged and older	67.2	2272	Self-report	8	1491	0.66	Moderate
¹⁶² Nunes et al (2016)	Brazil	South America	Low or middle	Community	All adults	45.75	2927	Self-report	11	852	0.29	Moderate
¹⁶³ Nunes et al (2017)	Brazil	South America	Low or middle	Community	All adults	43.7	60202	Self-report	22	13365	0.22	Moderate
¹⁶⁴ Nunes et al (2015)	Brazil	South America	Low or middle	Community	Only older people	Not reported	1593	Self-report	17	1295	0.81	Moderate
¹⁶⁵ Olaya et al (2017)	Spain	Europe	High	Community	Only older people	71.75	2113	Self-report	7	1088	0.51	Moderate
¹⁶⁶ Olivares et al (2017)	Argentina	South America	High	Community	All adults	43	1044	Self-report	Not reported	346	0.33	Moderate
¹⁶⁷ Park et al (2018)	South Korea	Asia	High	Community	Middle aged and older	62.7	5996	Self-report	25	1607	0.27	Moderate
¹⁶⁸ Patel et al (2006)	Mexico	South America	Low or middle	Community	Middle aged and older	73	7852	Self-report	5	1833	0.23	Moderate
¹⁶⁹ Pati et al (2016)	India	Asia	Low or middle	Community	All adults	44.96	103	Self-report	18	24	0.23	Moderate
¹⁷⁰ Pati et al (2019)	India	Asia	Low or middle	Primary care	All adults	44	1649	Self-report	21	567	0.34	Moderate
¹⁷¹ Payne et al (2013)	UK	Europe	High	Primary care	All adults	49	180815	Medical records and administrative database	40	54945	0.30	Moderate
¹⁷² Perez et al (2020)	Sweden	Europe	High	Community	Only older people	72.8	2596	Self-report	60	2213	0.85	Moderate
¹⁷³ Petersen et al (2019)	South Africa	Africa	Low or middle	Primary care	All adults	Not reported	2549	Self-report	Not reported	893	0.35	Moderate
¹⁷⁴ Pfortmueller et al (2013)	Switzerland	Europe	High	Hospitals	All adults	Median age: 28	3170	Medical records and administrative database	18	1183	0.37	High

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹⁷⁵ Pressley et al (1999)	USA	North America	High	Hospitals	Only older people	Not reported	5934	Medical records and administrative database	Not reported	3534	0.60	Moderate
¹⁷⁶ Prior et al (2016)	Denmark	Europe	High	Community	All adults	Not reported	118410	Self-report	39	33937	0.29	Moderate
¹⁷⁷ Ribeiro et al (2018)	Brazil	South America	High	Community	Only older people	70	820	Self-report	8	270	0.33	Moderate
¹⁷⁸ Ruel et al (2014)	Australia	Australasia	High	Community	All adults	50	1854	Self-report	8	585	0.32	Moderate
¹⁷⁹ Ruel et al (2014)	China	Asia	Low or middle	Community	All adults	49	1020	Self-report	11	346	0.34	Moderate
¹⁸⁰ Ryan et al (2018)	Ireland	Europe	High	Community	Middle aged and older	Not reported	4823	Self-report	16	2588	0.54	Moderate
¹⁸¹ Schmidt et al (2016)	Austria, Belgium, Denmark, France, Germany, Italy, the Netherlands, Spain, Sweden, and Switzerland	Europe	High	Community	Only older people	Not reported	56609	Self-report	11	13794	0.24	Moderate
¹⁸² Schotker et al (2016)	Germany	Europe	High	Primary care	Middle aged and older	Median age:70	2547	Medical records and administrative database	14	251	0.10	Moderate
¹⁸³ Seo et al (2017)	South Korea	Asia	High	Community	Middle aged and older	Not reported	156747	Self-report	15	42006	0.27	Moderate
¹⁸⁴ She et al (2019)	China	Asia	Low or middle	Hospitals	Only older people	68.9	1497	Self-report	22	1255	0.84	Moderate
¹⁸⁵ Singh et al (2019)	India	Asia	Low or middle	Community	All adults	41	16287	Self-report	5	1531	0.09	Moderate
¹⁸⁶ Stepanova et al (2015)	USA	North America	High	Community	All adults	34.7	26225	Self-report	13	9992	0.38	High
¹⁸⁷ Sticklely et al (2020)	USA	North America	High	Community	All adults	44.9	15311	Self-report	9	3996	0.26	High

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
¹⁸⁸ Streit et al (2014)	Switzerland	Europe	High	Primary care	Middle aged and older	63.5	1002	Medical records and administrative database	17	676	0.67	Moderate
¹⁸⁹ Stubbs et al (2018)	China, Ghana, India, Mexico, Russia, South Africa	Multiple continents	Low or middle	Community	Middle aged and older	62.4	34129	Self-report	13	19317	0.57	Moderate
¹⁹⁰ Su et al (2016)	China	Asia	Low or middle	Community	Only older people	Not reported	2058	Self-report	10	1012	0.49	Moderate
¹⁹¹ Sundstrup et al (2017)	USA	North America	High	Community	All adults	43.5	10427	Self-report	8	2489	0.24	High
¹⁹² Takahashi et al (2016)	USA	North America	High	Hospitals	All adults	57	6402	Medical records and administrative database	Not reported	3140	0.49	High
¹⁹³ Tinetti et al (2011)	USA	North America	High	Community	Only older people	72.6	5298	Self-report	5	1200	0.23	High
¹⁹⁴ Troelstra et al (2020)	The Netherlands	Europe	High	Community	All adults	Not reported	604	Self-report	26	321	0.53	High
¹⁹⁵ van Zon et al (2020)	USA	North America	High	Community	Middle aged and older	53.8	10719	Self-report	8	2390	0.22	Moderate
¹⁹⁶ Vancampfort et al (2017)	China, Ghana, India, Mexico, Russia, and South Africa	Multiple continents	Low or middle	Community	All adults	Median age: 62	32585	Self-report	11	14524	0.45	Moderate
¹⁹⁷ Vassilaki et al (2015)	USA	North America	High	Primary care	Only older people	78.5	2176	Medical records and administrative database	17	1884	0.87	Moderate
¹⁹⁸ Vassilaki et al (2016)	USA	North America	High	Primary care	Only older people	79	1449	Medical records and administrative database	17	1237	0.85	Moderate
¹⁹⁹ Villarreal et al (2015)	Panama	South America	High	Primary care	Only older people	78.2	304	Self-report	7	227	0.75	Moderate
²⁰⁰ Violan et al (2019)	Spain	Europe	High	Primary care	Only older people	75.4	916619	Medical records and	60	853085	0.93	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
								administrative database				
²⁰¹ von Strauss et al (2000)	Sweden	Europe	High	Community	Only older people	Not reported	502	Self-report	15	155	0.31	Moderate
²⁰² Vos et al (2013)	The Netherlands	Europe	High	Community	Only older people	71.9	315	Self-report	21	202	0.64	Moderate
²⁰³ Vu et al (2019)	Vietnam	Asia	Low or middle	Hospitals	Only older people	71.9	405	Medical records and administrative database	Not reported	146	0.36	High
²⁰⁴ Wang et al (2018)	USA	North America	High	Community	All adults	47	3086	Self-report	20	1109	0.36	Moderate
²⁰⁵ Wang et al (2017)	China	Asia	Low or middle	Community	Only older people	69.24	2705	Self-report	17	1230	0.45	Moderate
²⁰⁶ Wijers et al (2019)	Spain	Europe	High	Community	Middle aged and older	74.2	707	Self-report	21	491	0.69	Moderate
²⁰⁷ Williams et al (2016)	USA	North America	High	Community	All adults	Not reported	23789	Self-report	9	9213	0.39	Moderate
²⁰⁸ Woldeesemayat et al (2018)	Ethiopia	Africa	Low or middle	Primary care	All adults	Not reported	411	Self-report	18	73	0.18	Moderate
²⁰⁹ Yao et al (2020)	China	Asia	Low or middle	Community	Middle aged and older	57.7	10084	Self-report	15	3243	0.32	Moderate
²¹⁰ Yorke et al (2017)	USA	North America	High	Community	Middle aged and older	66.6	5877	Self-report	7	3391	0.58	Moderate
²¹¹ You et al (2019)	China	Asia	Low or middle	Community	Only older people	72	5296	Self-report	27	2201	0.42	Moderate
²¹² Zhang et al (2020)	China	Asia	Low or middle	Community	Only older people	74.14	4348	Self-report	15	2338	0.54	Moderate
²¹³ Khanam et al (2011)	Bangladesh	Asia	Low or middle	Community	Only older people	69.5	452	Medical records and administrative database	9	243	0.54	Moderate

Author	Country	Continent	Country income	Setting	Study population	Mean age	No of participants	Source	No of conditions measured	No of MM cases	Proportion with MM	Risk of bias
²¹⁴ Cornell et al (2009)	USA	North America	High	Primary care	All adults	62.4	1645314	Medical records and administrative database	45	1327382	0.81	Moderate
²¹⁵ Cassell et al (2018)	UK	Europe	High	Primary care	All adults	Not reported	403985	Medical records and administrative database	36	109884	0.27	Moderate
²¹⁶ Wong et al (2019)	Hong Kong (SAR of China)	Asia	High	Community	All adults	45.67	1014	Self-report	5	124	0.12	Moderate
²¹⁷ Puth et al (2017)	Germany	Europe	High	Community	All adults	Not reported	19294	Self-report	17	7640	0.40	Moderate

MM: Multimorbidity. No of participants is the total number of participants in the denominator for estimating prevalence in a study (which could be a subset in some included studies)

Table S5: Associations between predictors

	Mean age (lm) Unadjusted coefficient estimates	No of conditions (nb) Unadjusted incident rate ratio
Mean age		1.0 (1.0-1.0)
Source		
Self-report	59.7 (57.1-62.3) (intercept)	Ref
Database	7.0 (1.5-12.5)*	1.8 (1.5-2.2)***
Continent		
Europe	66.8 (62.8-70.9) (intercept)	Ref
North America	-7.0 (-12.8 to -1.1)*	0.6 (0.5-0.8)***
Australasia	-8.0 (-17.5-1.6)	0.8 (0.6-1.1)
Asia	-8.4 (-14.6 to -2.2)**	0.6 (0.5-0.8)***
South America	-8.5 (-18.0-1.1)	0.6 (0.4-0.9)**
Africa	-32.8 (-57.8 to -8.0)**	0.4 (0.2-0.8)*
Multiple continents	-7.6 (-18.3-3.2)	0.5 (0.3-0.7)***
Setting		
Community	59.8 (57.2-62.5) (intercept)	Ref
Primary care	3.5 (-2.5-9.6)	1.7 (1.4-2.1)***
Hospitals	10.2 (1.5-19.0)*	1.8 (1.3-2.4)***
Study population		
All adults	48.3 (46.6-50.0) (intercept)	Ref
Middle-aged and older	15.4 (12.7-18.0)***	0.9 (0.7-1.1)
Only older people	26.2 (23.7-28.7)***	1.2 (0.9-1.4)

* <0.05 ** <0.01 *** <0.001

Ref: Reference category. lm: Linear regression. nb: Negative binomial regression

Table S6: Risk of bias assessment of included studies

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
^{25.} Aarts et al (2012)	Moderate	Moderate	Moderate	High	Low	Low	Moderate	Low	Moderate	Yes
^{26.} Aarts et al (2011)	Low	High	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	No
^{27.} Aarts et al (2011)	Moderate	Moderate	Moderate	High	Low	Low	Moderate	Low	Moderate	Yes
^{28.} Abizanda et al (2014)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
^{29.} Agborsangaya et al (2012)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{30.} Agborsangaya et al (2013)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{31.} Agborsangaya et al (2014)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{32.} Ahrenfeldt et al (2019)	Low	Moderate	Moderate	High	Low	High	Moderate	Low	Moderate	No
^{33.} Alimohammadian et al (2017)	Low	Moderate	Moderate	High	Low	High	Moderate	Low	Moderate	Yes
^{34.} Angst et al (2002)	Moderate	Moderate	Moderate	High	Low	High	High	Unclear	High	No
^{35.} Appa et al (2014)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{36.} Adams et al (2017)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{37.} Ahmadi et al (2016)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{38.} Amaral et al (2018)	High	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
³⁹ . An et al (2016)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
⁴⁰ . Araujo et al (2018)	Low	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes
⁴¹ . Arnold-Reed et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁴² . Arokiasamy et al (2015)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁴³ . Sinnige et al (2015)	Low	Moderate	Moderate	High	Low	Low	Moderate	Low	Moderate	Yes
⁴⁴ . Zemedikun et al (2018)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁴⁵ . Wensing et al (2001)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Unclear	Moderate	Yes
⁴⁶ . Mounce et al (2018)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁴⁷ . Taylor et al (2010)	Low	Moderate	Moderate	High	Low	Low	Moderate	Low	Low	Yes
⁴⁸ . Vancampfort et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁴⁹ . Vancampfort et al (2018)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁵⁰ . Aubert et al (2016)	Moderate	Moderate	Moderate	High	Moderate	Low	Moderate	Low	Moderate	Yes
⁵¹ . Autenrieth et al (2013)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁵² . Bahler et al (2015)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Low	Moderate	Yes
⁵³ . Vancampfort et al (2017)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
⁵⁴ . Banjare et al (2014)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
⁵⁵ . Barra et al (2015)	Moderate	Moderate	High	High	Moderate	High	Moderate	Unclear	Moderate	No
⁵⁶ . Bernard et al (2016)	High	Moderate	High	High	Moderate	Low	Moderate	Low	High	No
⁵⁷ . Biswas et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
⁵⁸ . Blakemore et al (2016)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
⁵⁹ . Blyth et al (2008)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁶⁰ . Bowling et al (2019)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Low	Moderate	Yes
⁶¹ . Britt et al (2008)	Low	Moderate	High	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
⁶² . Broeiro-Goncalves (2019)	Low	Moderate	High	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
⁶³ . Bruce et al (2010)	High	Moderate	Moderate	High	Low	High	Moderate	Unclear	High	No
⁶⁴ . Burgers et al (2010)	Low	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	Yes
⁶⁵ . Burke et al (2017)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁶⁶ . Buurman et al (2016)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁶⁷ . Calderon-Larranaga et al (2017)	Moderate	Moderate	Moderate	High	Low	Low	Moderate	Low	Moderate	Yes
⁶⁸ . Camargo-Casas et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
^{69.} Canevelli et al (2019)	High	High	High	High	Moderate	High	Moderate	Low	High	Yes
^{70.} Chamberlain et al (2020)	Low	Moderate	Moderate	High	Low	Low	Moderate	Low	Low	Yes
^{71.} Chen et al (2018)	Low	Moderate	High	High	Low	Low	Moderate	Low	Low	Yes
^{72.} Chen et al (2018)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{73.} Cheung et al (2013)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{74.} Chu et al (2018)	Low	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
^{75.} Chudasama et al (2019)	Moderate	Moderate	Low	High	Moderate	Low	Moderate	Low	Moderate	Yes
^{76.} Cimarras-Otal et al (2014)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{77.} Chin et al (2016)	Moderate	Moderate	Moderate	High	Moderate	Low	Moderate	Low	Moderate	Yes
^{78.} Agrawal et al (2016)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{79.} Gu et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{80.} Gunn et al (2012)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{81.} Han et al (2013)	High	High	Moderate	High	Moderate	High	Moderate	Unclear	High	No
^{82.} Hanlon et al (2018)	Low	Moderate	Moderate	High	Low	Low	Moderate	Low	Low	Yes
^{83.} Jantsch et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
⁸⁴ John et al (2003)	Moderate	High	Moderate	High	Low	High	Moderate	Low	High	No
⁸⁵ Johnson-Lawrence et al (2017)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁸⁶ Johnston et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
⁸⁷ Jones et al (2016)	Low	Moderate	Moderate	High	Low	Low	Moderate	Unclear	Moderate	Yes
⁸⁸ Jovic et al (2016)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁸⁹ Juul-Larsen et al (2020)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁹⁰ Hudon et al (2008)	Low	Moderate	Moderate	High	Low	Low	Moderate	Low	Low	Yes
⁹¹ Hussain et al (2015)	Moderate	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	Yes
⁹² Ie et al (2017)	High	High	Moderate	High	Moderate	Low	Moderate	Low	High	Yes
⁹³ Ishizaki et al (2019)	Moderate	Moderate	Low	High	Moderate	High	Moderate	Unclear	Moderate	Yes
⁹⁴ Danon-Hersch et al (2012)	Moderate	High	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
⁹⁵ de Heer et al (2013)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
⁹⁶ Demirchyan et al (2013)	High	Moderate	Low	High	Moderate	High	Moderate	Low	High	No
⁹⁷ Fabbri et al (2015)	Moderate	Moderate	High	High	Moderate	High	Moderate	Unclear	Moderate	Yes
⁹⁸ Fillenbaum et al (2000)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
^{99.} Kaneko et al (2019)	Moderate	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	No
^{100.} Kang et al (2017)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{101.} Gandhi et al (2020)	Moderate	Moderate	Moderate	High	High	High	Moderate	Low	Moderate	Yes
^{102.} Costa et al (2018)	High	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{103.} Rizzuto et al (2017)	High	Moderate	High	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
^{104.} Dhalwani et al (2017)	Moderate	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes
^{105.} Elixhauser et al (1998)	Low	Moderate	High	High	Low	Low	Moderate	Unclear	Low	Yes
^{106.} Fabbri et al (2015)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{107.} Fortin et al (2014)	Low	Moderate	Low	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
^{108.} Fuchs et al (1998)	Moderate	Moderate	Moderate	High	Moderate	Low	Moderate	Unclear	Moderate	No
^{109.} Galenkamp et al (2011)	Low	Moderate	High	High	Moderate	High	Moderate	Unclear	High	No
^{110.} Galenkamp et al (2016)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{111.} Gamma et al (2001)	High	Moderate	High	High	Moderate	High	Moderate	Unclear	High	No
^{112.} Ge et al (2018)	Moderate	Moderate	High	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{113.} Ge et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
¹¹⁴ . Gould et al (2016)	Moderate	Moderate	Moderate	High	High	High	Moderate	Unclear	Moderate	Yes
¹¹⁵ . Habib et al (2014)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	No
¹¹⁶ . Harrison et al (2017)	Low	Moderate	High	High	Moderate	Low	Moderate	Unclear	Moderate	No
¹¹⁷ . Hayek et al (2017)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹¹⁸ . Henninger et al (2012)	Moderate	Moderate	High	High	Moderate	High	Moderate	Unclear	Moderate	No
¹¹⁹ . Hernandez et al (2019)	Moderate	Moderate	Moderate	High	High	High	Moderate	Unclear	Moderate	Yes
¹²⁰ . Ho et al (2014)	Moderate	Moderate	High	High	Low	Low	Moderate	Low	Moderate	Yes
¹²¹ . Khan et al (2019)	Low	Moderate	Low	High	Low	High	Moderate	Low	Low	Yes
¹²² . Kiliari et al (2013)	High	Moderate	High	High	Moderate	Moderate	Moderate	Low	Moderate	No
¹²³ . King et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹²⁴ . Kingston et al (2018)	Low	Moderate	High	High	Moderate	High	Moderate	Unclear	High	Yes
¹²⁵ . Koyanagi et al (2018)	Low	Moderate	Moderate	High	Moderate	Low	High	Low	Moderate	Yes
¹²⁶ . Kriegsman et al (2004)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹²⁷ . Kristensen et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹²⁸ . Kristensen et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
¹²⁹ . Kunna et al (2017)	Low	Moderate	Low	High	Moderate	Low	High	Low	Low	Yes
¹³⁰ . Kuwornu et al (2014)	Moderate	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	Yes
¹³¹ . Lai et al (2019)	Low	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	Yes
¹³² . Lai et al (2018)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹³³ . Laires et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹³⁴ . Lang et al (2015)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
¹³⁵ . Le Cossec et al (2016)	Low	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
¹³⁶ . Lee et al (2007)	Low	Moderate	High	High	Low	Low	Moderate	Low	Low	Yes
¹³⁷ . Lee et al (2018)	Low	Moderate	High	High	High	Low	Moderate	Unclear	High	No
¹³⁸ . Li et al (2016)	Low	Low	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹³⁹ . Li et al (2019)	Low	Moderate	Low	High	Moderate	Moderate	Moderate	Low	Moderate	No
¹⁴⁰ . Lujic et al (2017)	Low	Moderate	Moderate	High	Low	High	Moderate	Low	Moderate	Yes
¹⁴¹ . LupianezUnclearVillanueva et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹⁴² . Zhou et al (2018)	Moderate	Moderate	Moderate	High	Moderate	Low	High	Low	Moderate	Yes
¹⁴³ . Zhang et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
^{144.} Wong et al (2010)	High	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{145.} Weimann et al (2016)	Low	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes
^{146.} Wang et al (2017)	Low	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes
^{147.} Wang et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
^{148.} Wade et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{149.} Maciejewski et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{150.} Marengoni et al (2016)	Moderate	Moderate	High	High	Moderate	High	Moderate	Low	Moderate	Yes
^{151.} Marengoni et al (2009)	Moderate	Moderate	High	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{152.} Marques et al (2018)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{153.} Mavaddat et al (2014)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{154.} McDaid et al (2013)	Low	Moderate	High	High	Moderate	High	Moderate	Low	High	Yes
^{155.} Melis et al (2014)	High	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{156.} Min et al (2007)	High	Moderate	High	High	Moderate	High	Moderate	Unclear	High	Yes
^{157.} Momtaz et al (2010)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
^{158.} Mondor et al (2018)	Low	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
^{159.} Muggah et al (2012)	Low	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	No
^{160.} Nagel et al (2008)	Low	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes
^{161.} Niedzwiedz et al (2019)	Moderate	Moderate	High	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
^{162.} Nunes et al (2016)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{163.} Nunes et al (2017)	Low	Moderate	Moderate	High	Low	High	Moderate	Low	Moderate	Yes
^{164.} Nunes et al (2015)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{165.} Olaya et al (2017)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{166.} Olivares et al (2017)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
^{167.} Park et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
^{168.} Patel et al (2006)	Moderate	Moderate	High	High	Moderate	High	Moderate	Unclear	Moderate	No
^{169.} Pati et al (2016)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
^{170.} Pati et al (2019)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
^{171.} Payne et al (2013)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
^{172.} Perez et al (2020)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
^{173.} Petersen et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
¹⁷⁴ . Pfortmueller et al (2013)	Moderate	Moderate	High	High	High	High	Moderate	Unclear	High	No
¹⁷⁵ . Pressley et al (1999)	Low	Moderate	High	High	Moderate	Low	Moderate	Unclear	Moderate	No
¹⁷⁶ . Prior et al (2016)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
¹⁷⁷ . Ribeiro et al (2018)	High	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹⁷⁸ . Ruel et al (2014)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹⁷⁹ . Ruel et al (2014)	Moderate	Moderate	Moderate	High	Low	High	Moderate	Low	Moderate	Yes
¹⁸⁰ . Ryan et al (2018)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹⁸¹ . Schmidt et al (2016)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹⁸² . Schottker et al (2016)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes
¹⁸³ . Seo et al (2017)	Low	Moderate	Moderate	High	Low	High	Moderate	Low	Moderate	No
¹⁸⁴ . She et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹⁸⁵ . Singh et al (2019)	Low	Moderate	Moderate	High	Low	Low	Moderate	Unclear	Moderate	Yes
¹⁸⁶ . Stepanova et al (2015)	Low	High	High	High	High	High	High	Unclear	High	Yes
¹⁸⁷ . Stickley et al (2020)	Low	Moderate	High	High	Moderate	High	Moderate	Low	High	Yes
¹⁸⁸ . Streit et al (2014)	Moderate	Moderate	Moderate	High	High	High	Moderate	Unclear	Moderate	Yes

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
¹⁸⁹ . Stubbs et al (2018)	Low	Moderate	High	High	Moderate	Low	Moderate	Low	Moderate	Yes
¹⁹⁰ . Su et al (2016)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
¹⁹¹ . Sundstrup et al (2017)	Low	Moderate	High	High	Moderate	High	Moderate	Unclear	High	Yes
¹⁹² . Takahashi et al (2016)	Moderate	Moderate	High	High	High	Low	Moderate	Low	High	No
¹⁹³ . Tinetti et al (2011)	Low	Moderate	High	High	High	High	Moderate	Unclear	High	No
¹⁹⁴ . Troelstra et al (2020)	High	Moderate	High	High	Moderate	Low	Moderate	Unclear	High	Yes
¹⁹⁵ . van Zon et al (2020)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
¹⁹⁶ . Vancampfort et al (2017)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Low	Moderate	Yes
¹⁹⁷ . Vassilaki et al (2015)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
¹⁹⁸ . Vassilaki et al (2016)	Low	Moderate	Moderate	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
¹⁹⁹ . Villarreal et al (2015)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
²⁰⁰ . Violan et al (2019)	Low	Moderate	Moderate	High	High	Low	Moderate	Low	Moderate	Yes
²⁰¹ . von Strauss et al (2000)	Moderate	Moderate	Moderate	High	Moderate	Low	Moderate	Low	Moderate	No
²⁰² . Vos et al (2013)	Moderate	Moderate	High	High	Moderate	High	Moderate	Low	Moderate	No
²⁰³ . Vu et al (2019)	High	Moderate	High	High	Moderate	High	Moderate	Low	High	No

Author	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Publication bias	Conflict of interest	Overall rating	Reporting of MM measure and definition
²⁰⁴ . Wang et al (2018)	Moderate	Moderate	Low	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
²⁰⁵ . Wang et al (2017)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
²⁰⁶ . Wijers et al (2019)	Low	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	No
²⁰⁷ . Williams et al (2016)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	No
²⁰⁸ . Woldesemayat et al (2018)	High	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
²⁰⁹ . Yao et al (2020)	Moderate	Moderate	Low	High	Moderate	High	Moderate	Low	Moderate	Yes
²¹⁰ . Yorke et al (2017)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
²¹¹ . You et al (2019)	Moderate	Moderate	Moderate	High	Moderate	High	Moderate	Low	Moderate	Yes
²¹² . Zhang et al (2020)	Moderate	Moderate	Low	High	Moderate	Low	Moderate	Low	Moderate	Yes
²¹³ . Khanam et al (2011)	Moderate	Moderate	Moderate	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
²¹⁴ . Cornell et al (2009)	Low	Moderate	High	High	Moderate	Low	Moderate	Unclear	Moderate	Yes
²¹⁵ . Cassell et al (2018)	Low	Moderate	High	High	Moderate	Moderate	Moderate	Low	Moderate	No
²¹⁶ . Wong et al (2019)	High	Moderate	Moderate	High	Moderate	High	Moderate	Unclear	Moderate	Yes
²¹⁷ . Puth et al (2017)	Moderate	Moderate	Moderate	High	Moderate	Moderate	Moderate	Low	Moderate	Yes

Table S7: Output of adjusted meta-analytic model based on 217 studies

	Pooled prevalence of multimorbidity of each subgroup (% , 95% CI)	Meta-regression Unadjusted Odds Ratio (95% CI)	Meta-regression Adjusted Odds Ratio (95% CI) R ² 42.4%	FMI
Group of mean age		R ² 27.0%		
<59	30.4 (27.0-33.9)	Ref	Ref	Ref
59-73	43.5 (38.0-49.1)	1.8 (1.3-2.3)***	2.0 (1.6-2.6)***	0.3
≥74	67.8 (61.3-73.7)	6.4 (4.6-8.9)***	4.7 (3.4-6.5)***	0.2
No of conditions		R ² 6.9%		
<9	29.9 (24.9-35.4)	Ref	Ref	Ref
9-19	43.5 (39.1-47.9)	1.8 (1.3-2.5)***	1.7 (1.3-2.2)***	0.1
20-43	46.7 (38.4-55.2)	2.1 (1.4-3.1)***	2.2 (1.5-3.3)***	0.2
≥44	54.5 (32.6-74.8)	2.8 (1.5-5.4)**	2.8 (1.6-4.8)***	0.1
Setting		R ² 3.7%		
Community	37.8 (34.4-41.4)	Ref	Ref	Ref
Primary care	51.2 (41.6-60.7)	1.7 (1.2-2.5)**	1.8 (1.2-2.6)**	0.1
Hospital	47.1 (31.9-63.0)	1.5 (0.9-2.4)	0.8 (0.5-1.3)	0.1
Care home	73.9 (72.8-74.9)	4.6 (0.6-36.6)	1.5 (0.3-8.4)	0.04
Source		R ² 2.8%		
Self-report	38.3 (34.4-42.2)	Ref	Ref	Ref
Database	48.9 (42.2-55.6)	1.5 (1.1-2.1)**	0.8 (0.6-1.1)	0.1
Continent		R ² 7.4%		
North America	48.9 (42.1-55.7)	Ref	Ref	Ref
Europe	44.0 (37.7-50.4)	0.8 (0.6-1.2)	0.5 (0.4-0.7)***	0.1
Australasia	28.2 (20.3-37.6)	0.4 (0.2-0.8)**	0.4 (0.2-0.6)***	0.08
Asia	34.3 (28.6-40.5)	0.5 (0.4-0.8)**	0.5 (0.3-0.7)***	0.1
South America	47.5 (31.2-64.4)	0.9 (0.5-1.8)	0.8 (0.5-1.3)	0.1
Africa	13.8 (4.5-35.2)	0.2 (0.06-0.5)***	0.2 (0.1-0.5)***	0.1
Multiple continents	41.4 (31.0-52.6)	0.7 (0.4-1.4)	0.7 (0.4-1.2)	0.1


* <0.05 ** <0.01 *** <0.001

Ref: Reference category. FMI: Fraction of missing information.

Table S8: Definition of variables

Variable name	Definition
Study setting	
Community	Studies that used population surveys, insurance claims databases, or research databases
Primary care	Studies that were carried out in primary care settings
Hospital	Studies that were carried out in hospital settings
Data source	
Self-report	Studies that collected data using self-report or interviews
Medical records and administrative databases	Studies that collected data using electronic medical records, medical chart reviews, insurance claims databases, pharmacy databases, or research databases
Study population	
All adults	Studies with a sample of population aged 18 and older (n=45), aged 20 and older (n=8), aged 21 and older (n=3), aged 25 and older (n=2), or others (n=27) (e.g. aged 16 and older, or aged 17 and older)
Middle-aged and older	Studies with a sample of population aged 50 and older (n=25), aged 40 and older (n=5), aged 40 and older (n=10), or others (n=6) (e.g. aged 57 and older, or aged 45 and older)
Only older people	Studies with a sample of population aged 65 and older (n=22), aged 60 and older (n=25), aged 70 and older (n=5) or others (n=11) (e.g. aged 68 and older, aged 77 and older, aged 78 and older, or aged 80 and older)

Figure S1: Graphical display of study effect sizes and heterogeneity



No obvious subgroup effects were identified

Figure S2: Process of examining and identifying outlying studies in meta-analysis






Figure S3: Summary of risk of bias assessment

Figure S4: Meta-regression trees for predicting the pooled estimated prevalence of multimorbidity (based on ‘mean age’ and ‘number of conditions’ predictors. unit: log(odds))



References

- 1 Stanley J, Semper K, Millar E, Sarfati D. Epidemiology of multimorbidity in New Zealand: A cross-sectional study using national-level hospital and pharmaceutical data. *BMJ Open* 2018; **8**:1-12.
- 2 Lenzi J, Avaldi VM, Rucci P, Pieri G, Fantini MP. Burden of multimorbidity in relation to age, gender and immigrant status: A cross-sectional study based on administrative data. *BMJ Open* 2016; **6**:1-12.
- 3 Hu RH, Hsiao FY, Chen LJ, Huang PT, Hsu WW. Increasing age- and gender-specific burden and complexity of multimorbidity in taiwan, 2003-2013: A cross-sectional study based on nationwide claims data. *BMJ Open* 2019; **9**:1-10.
- 4 Gawron LM, Sanders JN, Sward K, Poursaid AE, Simmons R, Turok DK. Multimorbidity and highly effective contraception in reproductive-age women in the us intermountain west: A retrospective cohort study. *J Gen Intern Med* 2020; **35**:637-42.
- 5 Low LL, Kwan YH, Ko MSM, et al. Epidemiologic characteristics of multimorbidity and sociodemographic factors associated with multimorbidity in a rapidly aging asian country. *JAMA Network Open* 2019; **2**:1-14.
- 6 Wang HHX, Wang JJ, Wong SYS, et al. Epidemiology of multimorbidity in china and implications for the healthcare system: Cross-sectional survey among 162,464 community household residents in southern china. *BMC Medicine* 2014; **12**:1-12.
- 7 Gaulin M, Simard M, Candas B, Lesage A, Sirois C. Combined impacts of multimorbidity and mental disorders on frequent emergency department visits: A retrospective cohort study in quebec, canada. *Canadian Medical Association Journal* 2019; **191**:E724-E32.
- 8 Violan C, Foguet-Boreu Q, Flores-Mateo G, et al. Prevalence, determinants and patterns of multimorbidity in primary care: A systematic review of observational studies. *PLOS One* 2014; **9**:1-9.
- 9 Nicholson K, Terry AL, Fortin M, Williamson T, Bauer M, Thind A. Prevalence, characteristics, and patterns of patients with multimorbidity in primary care: A retrospective cohort analysis in canada. *British Journal of General Practice* 2019; **69**:1-10.
- 10 Bao X-Y, Xie Y-X, Zhang X-X, et al. The association between multimorbidity and health-related quality of life: A cross-sectional survey among community middle-aged and elderly residents in southern china. *Health and quality of life outcomes* 2019; **17**:107-09.
- 11 Fortin M, Bravo G, Hudon C, Vanasse A, Lapointe L. Prevalence of multimorbidity among adults seen in family practice. *Ann Fam Med* 2005; **3**:223-8.
- 12 Prazeres F Santiago L. Prevalence of multimorbidity in the adult population attending primary care in portugal: A cross-sectional study. *BMJ Open* 2015; **5**:1-10.
- 13 Lawson KD, Mercer SW, Wyke S, et al. Double trouble: The impact of multimorbidity and deprivation on preference-weighted health related quality of life a cross sectional analysis of the scottish health survey. *Int J Equity Health* 2013; **12**:1-9.
- 14 Sullivan PW, Ghushchyan VH, Bayliss EA. The impact of co-morbidity burden on preference-based health-related quality of life in the United States. *Pharmacoeconomics* 2012; **30**:431-42.
- 15 Peng X, Bao X, Xie Y, et al. The mediating effect of pain on the association between multimorbidity and disability and impaired physical performance among community-dwelling older adults in southern china. *Aging Clin Exp Res* 2020; **32**:1327-34.

- 16 Excoffier S, Herzig L, N'Goran AA, Déruaz-Luyet A, Haller DM. Prevalence of multimorbidity in general practice: A cross-sectional study within the Swiss sentinel surveillance system (sentinella). *BMJ Open* 2018; **8**:1-7.
- 17 Chung RY, Mercer S, Lai FTT, Yip BHK, Wong MCS, Wong SYS. Socioeconomic determinants of multimorbidity: A population-based household survey of Hong Kong Chinese. *PLOS ONE* 2015; **10**:1-15.
- 18 Ki M, Lee YH, Kim Y-S, Shin J-Y, Lim J, Nazroo J. Socioeconomic inequalities in health in the context of multimorbidity: A Korean panel study. *PLOS ONE* 2017; **12**:1-14.
- 19 Bobo WV, Yawn BP, St Sauver JL, Grossardt BR, Boyd CM, Rocca WA. Prevalence of combined somatic and mental health multimorbidity: Patterns by age, sex, and race/ethnicity. *J Gerontol - A Biol Sci Med Sci* 2016; **71**:1483-91.
- 20 Randall DA, Lujic S, Havard A, Eades SJ, Jorm L. Multimorbidity among aboriginal people in New South Wales contributes significantly to their higher mortality. *Med J Aust* 2018; **209**:19-23.
- 21 Russell J, Grant CC, Morton SMB. Multimorbidity in early childhood and socioeconomic disadvantage: Findings from a large New Zealand child cohort. *Acad Pediatr* 2020; **20**:619-27.
- 22 Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: A cross-sectional study. *The Lancet* 2012; **380**:37-43.
- 23 St Sauver JL, Boyd CM, Grossardt BR, et al. Risk of developing multimorbidity across all ages in an historical cohort study: Differences by sex and ethnicity. *BMJ Open* 2015; **5**:1-13.
- 24 Vetrano DL, Foebel AD, Marengoni A, et al. Chronic diseases and geriatric syndromes: The different weight of comorbidity. *Eur J Intern Med* 2016; **27**:62-7.
- 25 Aarts S, den Akker Mv, Bosma H, et al. The effect of multimorbidity on health related functioning: Temporary or persistent? Results from a longitudinal cohort study. *J Psychosom Res* 2012; **73**:211-17.
- 26 Aarts S, van den Akker M, Hajema KJ, et al. Multimorbidity and its relation to subjective memory complaints in a large general population of older adults. *Int Psychogeriatr* 2011; **23**:616-24.
- 27 Aarts S, van den Akker M, Tan FES, Verhey FRJ, Metsemakers JFM, van Boxtel MPJ. Influence of multimorbidity on cognition in a normal aging population: A 12-year follow-up in the maastricht aging study. *Int J Geriatr Psychiatry* 2011; **26**:1046-53.
- 28 Abizanda P, Romero L, Sanchez-Jurado PM, Martinez-Reig M, Alfonso-Silguero SA, Rodriguez-Manas L. Age, frailty, disability, institutionalization, multimorbidity or comorbidity. Which are the main targets in older adults? *J Nutr, Health Aging* 2014; **18**:622-27.
- 29 Agborsangaya CB, Lau D, Lahtinen M, Cooke T, Johnson JA. Health-related quality of life and healthcare utilization in multimorbidity: Results of a cross-sectional survey. *Qual Life Res* 2012; **22**:791-99.
- 30 Agborsangaya CB, Ngwakongwi E, Lahtinen M, Cooke T, Johnson JA. Multimorbidity prevalence in the general population: The role of obesity in chronic disease clustering. *BMC Public Health* 2013; **13**:1161-61.
- 31 Agborsangaya CB, Lahtinen M, Cooke T, Johnson JA. Comparing the eq-5d 3l and 5l: Measurement properties and association with chronic conditions and multimorbidity in the general population. *Health Qual Life Outcomes* 2014; **12**:1-7.

- 32 Ahrenfeldt LJ, Möller S, Thinggaard M, Christensen K, Lindahl-Jacobsen R. Sex differences in comorbidity and frailty in Europe. *Int J Public Health* 2019; **64**:1025-36.
- 33 Alimohammadian M, Majidi A, Yaseri M, et al. Multimorbidity as an important issue among women: Results of a gender difference investigation in a large population-based cross-sectional study in west asia. *BMJ Open* 2017; **7**:1-8.
- 34 Angst J, Sellaro R, Ries Merikangas K. Multimorbidity of psychiatric disorders as an indicator of clinical severity. *Eur Arch Psychiatry Clin Neurosci* 2002; **252**:147-54.
- 35 Appa AA, Creasman J, Brown JS, et al. The impact of multimorbidity on sexual function in middle-aged and older women: Beyond the single disease perspective. *J Sex Med* 2014; **11**:2744-55.
- 36 Adams ML, Grandpre J, Katz DL, Shenson D. Linear association between number of modifiable risk factors and multiple chronic conditions: Results from the behavioral risk factor surveillance system. *Prev Med* 2017; **105**:169-75.
- 37 Ahmadi B, Alimohammadian M, Yaseri M, et al. Multimorbidity: Epidemiology and risk factors in the golesstan cohort study, iran a cross-sectional analysis. *Medicine* 2016; **95**:1-7.
- 38 Amaral TLM, Amaral CdA, de Lima NS, Herculano PV, Do Prado PR, Monteiro GTR. Multimorbidity, depression and quality of life among elderly people assisted in the family health strategy in senador guiomard, acre, Brazil. *Ciencia e Saude Coletiva* 2018; **23**:3077-84.
- 39 An KO Kim J. Association of sarcopenia and obesity with multimorbidity in Korean adults: A nationwide cross-sectional study. *J Am Med Dir Assoc* 2016; **17**:e1-7.
- 40 Araujo MEA, Silva MT, Galvao TF, Nunes BP, Pereira MG. Prevalence and patterns of multimorbidity in amazon region of Brazil and associated determinants: A cross-sectional study. *BMJ Open* 2018; **8**:1-11.
- 41 Arnold-Reed D, Troeung L, Brett T, et al. Increasing multimorbidity in an Australian street health service: A 10-year retrospective cohort study. *Aust J Gen Pract* 2018; **47**:181-89.
- 42 Arokiasamy P, Uttamacharya U, Jain K, et al. The impact of multimorbidity on adult physical and mental health in low- and middle-income countries: What does the study on global ageing and adult health (sage) reveal? *BMC Med* 2015; **13**:1-16.
- 43 Sinnige J, Korevaar JC, Westert GP, Spreuwenberg P, Schellevis FG, Braspenning JC. Multimorbidity patterns in a primary care population aged 55 years and over. *Fam Pract* 2015; **32**:505-13.
- 44 Zemedikun DT, Gray LJ, Khunti K, Davies MJ, Dhalwani NN. Patterns of multimorbidity in middle-aged and older adults: An analysis of the uk biobank data. *Mayo Clin Proc* 2018; **93**:857-66.
- 45 Wensing M, Vingerhoets E, Grol R. Functional status, health problems, age and comorbidity in primary care patients. *Qual Life Res* 2001; **10**:141-48.
- 46 Mounce LTAP, Campbell JLMD, Henley WEP, Arreal MCTPMPHLCP, Porter IP, Valderas JMBPMPH. Predicting incident multimorbidity. *Ann Fam Med* 2018; **16**:322-29.
- 47 Taylor AW, Price K, Gill TK, et al. Multimorbidity - not just an older person's issue. Results from an Australian biomedical study. *BMC Public Health* 2010; **10**:1-10.
- 48 Vancampfort D, Stubbs B, Firth J, Koyanagi A. Handgrip strength, chronic physical conditions and physical multimorbidity in middle-aged and older adults in six low- and middle income countries. *Eur J Intern Med* 2019; **61**:96-102.

- 49 Vancampfort D, Smith L, Stubbs B, et al. Associations between active travel and physical multi-morbidity in six low- and middle-income countries among community-dwelling older adults: A cross-sectional study. *PloS one* 2018; **13**:1-15.
- 50 Aubert CE, Streit S, Da Costa BR, et al. Polypharmacy and specific comorbidities in university primary care settings. *Eur J Intern Med* 2016; **35**:35-42.
- 51 Autenrieth CS, Kirchberger I, Heier M, et al. Physical activity is inversely associated with multimorbidity in elderly men: Results from the kora-age augsburg study. *Prev Med* 2013; **57**:17-19.
- 52 Bähler C, Huber CA, Brüngger B, Reich O. Multimorbidity, health care utilization and costs in an elderly community-dwelling population: A claims data based observational study. *BMC Health Services Res* 2015; **15**:1-12.
- 53 Vancampfort DK, A.: Ward, P. B.: Veronese, N.: Carvalho, A. F.: Solmi, M.: Mugisha, J.: Rosenbaum, S.: De Hert, M.: Stubbs, B. Perceived stress and its relationship with chronic medical conditions and multimorbidity among 229,293 community-dwelling adults in 44 low- and middle-income countries. *Am J Epidemiol* 2017; **186**:979-89.
- 54 Banjare P Pradhan J. Socio-economic inequalities in the prevalence of multi-morbidity among the rural elderly in Bargarh district of Odisha (India). *PLoS ONE* 2014; **9**:1-10.
- 55 Barra M, Augestad LA, Whitehurst DGT, Rand-Hendriksen K. Examining the relationship between health-related quality of life and increasing numbers of diagnoses. *Qual life Res* 2015; **24**:2823-32.
- 56 Bernard S, Inderjeeth C, Raymond W. Higher charlson comorbidity index scores do not influence functional independence measure score gains in older rehabilitation patients. *Aust J Ageing* 2016; **35**:236-41.
- 57 Biswas T, Townsend N, Islam MS, et al. Association between socioeconomic status and prevalence of non-communicable diseases risk factors and comorbidities in bangladesh: Findings from a nationwide cross-sectional survey. *BMJ Open* 2019; **9**:1-9.
- 58 Blakemore A, Hann M, Howells K, et al. Patient activation in older people with long-term conditions and multimorbidity: Correlates and change in a cohort study in the united kingdom. *BMC Health Services Res* 2016; **16**:582-82.
- 59 Blyth FM, Rochat S, Cumming RG, et al. Pain, frailty and comorbidity on older men: The champ study. *Pain* 2008; **140**:224-30.
- 60 Bowling CB, Deng L, Sakhuja S, Morey MC, Jaeger BC, Muntner P. Prevalence of activity limitations and association with multimorbidity among us adults 50 to 64 years old. *J Gen Intern Med* 2019; **34**:2390-96.
- 61 Britt HC, Harrison CM, Miller GC, Knox SA. Prevalence and patterns of multimorbidity in australia. *Med J Aust* 2008; **189**:72-77.
- 62 Broeiro-Gonçalves P, Nogueira P, Aguiar P. Multimorbidity and disease severity by age groups, in inpatients: Cross-sectional study. *Portuguese J Public Health* 2019; **37**:1-9.
- 63 Bruce SG, Riediger ND, Zacharias JM, Young TK. Obesity and obesity-related comorbidities in a canadian first nation population. *Chronic Dis Can* 2010; **31**:27-32.
- 64 Burgers JS, Voerman GE, Grol R, Faber MJ, Schneider EC. Quality and coordination of care for patients with multiple conditions: Results from an international survey of patient experience. *Eval Health Prof* 2010; **33**:343-64.
- 65 Burke BT. *Connections that count: A network analysis of multimorbidity and gait speed in older adults*, in *Department of Epidemiology (dissertation)*. 2017, Johns Hopkins University: Baltimore, Maryland.1-224.

- 66 Buurman BM, Frenkel WJ, Abu-Hanna A, Parlevliet JL, de Rooij SE. Acute and chronic diseases as part of multimorbidity in acutely hospitalized older patients. *Eur J Intern Med* 2016; **27**:68-75.
- 67 Calderón-Larrañaga A, Vetrano DL, Onder G, et al. Assessing and measuring chronic multimorbidity in the older population: A proposal for its operationalization. *J Gerontol A Biol Sci Med Sci* 2017; **72**:1417-23.
- 68 Camargo-Casas S, Suarez-Monsalve S, Zepeda MUP, García-Peña C, Cano-Gutierrez CA. Multimorbidity, depressive symptoms, and self-reported health in older adults: A secondary analysis of the sabe bogota study. *Rev Invest Clin* 2018; **70**:192-97.
- 69 Canevelli M, Raganato R, Remiddi F, et al. Counting deficits or diseases? The agreement between frailty and multimorbidity in subjects with cognitive disturbances. *Aging Clin Exp Res* 2019; **32**:179-82.
- 70 Chamberlain AM, Rutten LJJ, Wilson PM, et al. Neighborhood socioeconomic disadvantage is associated with multimorbidity in a geographically-defined community. *BMC Public Health* 2020; **20**:1-10.
- 71 Chen H, Chen Y, Cui B. The association of multimorbidity with healthcare expenditure among the elderly patients in Beijing, China. *Arch Gerontol Geriatr* 2018; **79**:32-38.
- 72 Chen H, Cheng M, Zhuang Y, Broad JBJG, International G. Multimorbidity among middle-aged and older persons in urban china: Prevalence, characteristics and health service utilization. *Geriatr Gerontol Int* 2018; **18**:1447-52.
- 73 Cheung CL, Nguyen US, Au E, Tan KC, Kung AW. Association of handgrip strength with chronic diseases and multimorbidity: A cross-sectional study. *Age (Dordr)* 2013; **35**:929-41.
- 74 Chu T, Lau P, Cheng R, Chan M, Liang J. A study on the prevalence of multimorbidities of diseases and utilisation of public healthcare services in the new territories west area of Hong Kong. *Hong Kong Practitioner* 2018; **40**:43-50.
- 75 Chudasama YV, Khunti KK, Zaccardi F, et al. Physical activity, multimorbidity, and life expectancy: A UK biobank longitudinal study. *BMC Med* 2019; **17**:1-13.
- 76 Cimarras-Otal C, Calderón-Larrañaga A, Poblador-Plou B, et al. Association between physical activity, multimorbidity, self-rated health and functional limitation in the Spanish population. *BMC Public Health* 2014; **14**:1-10.
- 77 Chin WY, Choi EP, Wan EY, Lam CL. Health-related quality of life mediates associations between multi-morbidity and depressive symptoms in Chinese primary care patients. *Fam Pract* 2016; **33**:61-8.
- 78 Agrawal S, Agrawal PK. Association between body mass index and prevalence of multimorbidity in low-and middle-income countries: A cross-sectional study. *Int J Med Public Health* 2016; **6**:73-83.
- 79 Gu J, Chao J, Chen W, et al. Multimorbidity and health-related quality of life among the community-dwelling elderly: A longitudinal study. *Arch Gerontol Geriatr* 2018; **74**:133-40.
- 80 Gunn JM, Ayton DR, Densley K, et al. The association between chronic illness, multimorbidity and depressive symptoms in an australian primary care cohort. *Soc Psychiatry Psychiatr Epidemiol* 2010; **47**:175-84.
- 81 Han MA, Tinetti ME, Agostini JV, Han L, Lee H, Walke LM. The impact of medication use on symptom burden in older patients with multiple medical morbidities. *J Clin Gerontol and Geriatr* 2013; **4**:84-88.
- 82 Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality:

- A prospective analysis of 493 737 UK biobank participants. *Lancet Public Health* 2018; **3**:e323-e32.
- 83 Jantsch AG, Alves RFS, Faerstein E. Educational inequality in Rio de Janeiro and its impact on multimorbidity: Evidence from the pró-saúde study. A cross-sectional analysis. *Sao Paulo Med J* 2018; **136**:51-58.
- 84 John R, Kerby DS, Hagan Hennessy C. Patterns and impact of comorbidity and multimorbidity among community-resident American Indian elders. *Gerontol* 2003; **43**:649-60.
- 85 Johnson-Lawrence V, Zajacova A, Sneed R. Education, race/ethnicity, and multimorbidity among adults aged 30-64 in the national health interview survey. *SSM Popul Health* 2017; **3**:366-72.
- 86 Johnston MC, Black C, Mercer SW, Prescott GJ, Crilly MA. Impact of educational attainment on the association between social class at birth and multimorbidity in middle age in the Aberdeen children of the 1950s cohort study. *BMJ Open* 2019; **9**:1-9.
- 87 Jones SM, Amtmann D, Gell NM. A psychometric examination of multimorbidity and mental health in older adults. *Aging Ment Health* 2016; **20**:309-17.
- 88 Jovic D, Marinkovic J, Vukovic D. Association between body mass index and prevalence of multimorbidity: A cross-sectional study. *Public Health* 2016; **139**:103-11.
- 89 Juul-Larsen HG, Andersen O, Bandholm T, et al. Differences in function and recovery profiles between patterns of multimorbidity among older medical patients the first year after an acute admission—an exploratory latent class analysis. *Arch Gerontol Geriatr* 2020; **86**:1-11.
- 90 Hudon C, Soubhi H, Fortin M. Relationship between multimorbidity and physical activity: Secondary analysis from the Quebec health survey. *BMC Public Health* 2008; **8**:1-8.
- 91 Hussain MA, Huxley RR, Al Mamun A. Multimorbidity prevalence and pattern in Indonesian adults: An exploratory study using national survey data. *BMJ Open* 2015; **5**:1-10.
- 92 Ie K, Felton M, Springer S, Wilson SA, Albert SM. Multimorbidity and polypharmacy in family medicine residency practices. *J Pharm Technol* 2017; **33**:219-24.
- 93 Ishizaki T, Kobayashi E, Fukaya T, Takahashi Y, Shinkai S, Liang J. Association of physical performance and self-rated health with multimorbidity among older adults: Results from a nationwide survey in Japan. *Arch Gerontol Geriatr* 2019; **84**:1-9.
- 94 Danon-Hersch N, Rodondi N, Spagnoli J, Santos-Eggimann B. Prefrailty and chronic morbidity in the youngest old: An insight from the Lausanne cohort Lc65+. *J Am Geriatr Soc* 2012; **60**:1687-94.
- 95 de Heer HD, Balcázar HG, Morera OF, et al. Barriers to care and comorbidities along the U.S.-Mexico border. *Public Health Rep* 2013; **128**:480-8.
- 96 Demirchyan A, Khachadourian V, Armenian HK, Petrosyan V. Short and long term determinants of incident multimorbidity in a cohort of 1988 earthquake survivors in Armenia. *Int. J. Equity Health* 2013; **12**:68-68.
- 97 Fabbri E, An Y, Zoli M, et al. Aging and the burden of multimorbidity: Associations with inflammatory and anabolic hormonal biomarkers. *J Gerontol A Biol Sci Med Sci* 2015; **70**:63-70.
- 98 Fillenbaum GGP, C. F.: Cohen, H. J.: Cornoni-Huntley, J. C.: Guralnik, J. M. Comorbidity of five chronic health conditions in elderly community residents: Determinants and impact on mortality. *J Gerontol: Med Sci* 2000; **55**:M84-89.

- 99 Kaneko M, Boven Kv, Takayanagi H, Kusaba T, Yamada T, Matsushima MJFp. Multicentre descriptive cross-sectional study of japanese home visit patients: Reasons for encounter, health problems and multimorbidity. *Fam Pract* 2019; **37**:227-33.
- 100 Kang J Song Y-M. The association between submarine service and multimorbidity: A cross-sectional study of korean naval personnel. *BMJ Open* 2017; **7**:1-6.
- 101 Gandhi K, Lim E, Davis J, Chen JJ. Racial-ethnic disparities in self-reported health status among us adults adjusted for sociodemographics and multimorbidities, national health and nutrition examination survey 2011-2014. *Ethn Health* 2020; **25**:65-78.
- 102 Costa CDS, Flores TR, Wendt A, et al. Inequalities in multimorbidity among elderly: A population-based study in a city in southern brazil. *Cad Saude Publica* 2018; **34**:1-12.
- 103 Rizzuto DM, R. J. F.: Angleman, S.: Qiu, C.: Marengoni, A. Effect of chronic diseases and multimorbidity on survival and functioning in elderly adults. *J Am Geriatr Soc* 2017; **65**:1056-60.
- 104 Dhalwani NN, Zaccardi F, O'Donovan G, et al. Association between lifestyle factors and the incidence of multimorbidity in an older english population. *J Gerontol A Biol Sci Med Sci* 2017; **72**:528-34.
- 105 Elixhauser A, Steiner C, Harris DR, Coffey RM. Comorbidity measures for use with administrative data. *Med Care* 1998; **36**:8-27.
- 106 Fabbri E, An Y, Schrack JA, et al. Energy metabolism and the burden of multimorbidity in older adults: Results from the baltimore longitudinal study of aging. *J Gerontol A Biol Sci Med Sci* 2015; **70**:1297-303.
- 107 Fortin M, Haggerty J, Almirall J, Bouhali T, Sasseville M, Lemieux M. Lifestyle factors and multimorbidity: A cross sectional study. *BMC Public Health* 2014; **14**:1-6.
- 108 Fuchs ZB, T.: Novikov, I.: Walter-Ginzburg, A.: Lyanders, M.: Gindin, J.: Habot, B.: Modan, B. Morbidity, comorbidity, and their association with disability among community-dwelling oldest-old in israel. *J Gerontol: Med Sci* 1998; **53**:M447-55.
- 109 Galenkamp H, Braam AW, Huisman M, Deeg DJ. Somatic multimorbidity and self-rated health in the older population. *J Gerontol B Psychol Sci Soc Sci* 2011; **66**:380-6.
- 110 Galenkamp H, Gagliardi C, Principi A, et al. Predictors of social leisure activities in older europeans with and without multimorbidity. *Eur J Ageing* 2016; **13**:129-43.
- 111 Gamma A Angst J. Concurrent psychiatric comorbidity and multimorbidity in a community study: Gender differences and quality of life. *Eur Arch Psychiatry Clin Neurosci* 2001; **251**:43-46.
- 112 Ge L, Yap CW, Heng BH. Sex differences in associations between multimorbidity and physical function domains among community-dwelling adults in Singapore. *PLoS One* 2018; **13**:1-14.
- 113 Ge L, Ong R, Yap CW, Heng BH. Effects of chronic diseases on health-related quality of life and self-rated health among three adult age groups. *Nurs Health Sci* 2019; **21**:214-22.
- 114 Gould CE, O'Hara R, Goldstein MK, Beaudreau SA. Multimorbidity is associated with anxiety in older adults in the health and retirement study. *Int J Geriatr Psychiatry* 2016; **31**:1105-15.
- 115 Habib RR, Hojeij S, Elzein K, Chaaban J, Seyfert K. Associations between life conditions and multi-morbidity in marginalized populations: The case of palestinian refugees. *Eur J Public Health* 2014; **24**:727-33.
- 116 Harrison C, Henderson J, Miller G, Britt H. The prevalence of diagnosed chronic conditions and multimorbidity in australia: A method for estimating population prevalence from general practice patient encounter data. *PLOS ONE* 2017; **12**:1-13.

- 117 Hayek S, Ifrah A, Enav T, Shohat T. Prevalence, correlates, and time trends of multiple chronic conditions among israeli adults: Estimates from the Israeli national health interview survey, 2014-2015. *Prev Chronic Dis* 2017; **14**:1-14.
- 118 Henninger DE, Whitson HE, Cohen HJ, Ariely D. Higher medical morbidity burden is associated with external locus of control. *J Am Geriatr Soc* 2012; **60**:751-55.
- 119 Hernández B, Reilly RB, Kenny RA. Investigation of multimorbidity and prevalent disease combinations in older irish adults using network analysis and association rules. *Sci Rep* 2019; **9**:1-12.
- 120 Ho C, Feng L, Fam J, Mahendran R, Kua EH, Ng TP. Coexisting medical comorbidity and depression: Multiplicative effects on health outcomes in older adults. *Int Psychogeriatr* 2014; **26**:1221-9.
- 121 Khan N, Rahman M, Mitra D, Afsana K. Prevalence of multimorbidity among bangladeshi adult population: A nationwide cross-sectional study. *BMJ Open* 2019; **9**:1-8.
- 122 Kiliari N, Theodosopoulou E, Papanastasiou E. Multimorbidity and unmet citizens' needs and expectations urge for reforms in the health system of cyprus: A questionnaire survey. *JRSM Open* 2014; **5**:1-11.
- 123 King DE, Xiang J, Pilkerton CS. Multimorbidity trends in united states adults, 1988-2014. *J Am Board Fam Med* 2018; **31**:503-13.
- 124 Kingston AR, L.: Booth, H.: Knapp, M.: Jagger, C.: Modem project. Projections of multi-morbidity in the older population in england to 2035: Estimates from the population ageing and care simulation (pacsim) model. *Age Ageing* 2018; **47**:374-80.
- 125 Koyanagi A, Lara E, Stubbs B, et al. Chronic physical conditions, multimorbidity, and mild cognitive impairment in low- and middle-income countries. *J Am Geriatr Soc* 2018; **66**:721-27.
- 126 Kriegsman DMD, D. J.: Stalman, W. A. Comorbidity of somatic chronic diseases and decline in physical functioning;; the longitudinal aging study amsterdam. *J Clin Epidemiol* 2004; **57**:55-65.
- 127 Kristensen K, König HH, Hajek A. The longitudinal association of multimorbidity on loneliness and network size: Findings from a population-based study. *Int J Geriatr Psychiatry* 2019; **34**:1490-97.
- 128 Kristensen K, König HH, Hajek A. The association of multimorbidity, loneliness, social exclusion and network size: Findings from the population-based german ageing survey. *BMC Public Health* 2019; **19**:1-10.
- 129 Kunna R, San Sebastian M, Stewart Williams J. Measurement and decomposition of socioeconomic inequality in single and multimorbidity in older adults in china and ghana: Results from the who study on global ageing and adult health (sage). *Int J Equity Health* 2017; **16**:2-17.
- 130 Kusumastuti S, Gerds TA, Lund R, Mortensen EL, Westendorp RGJ. Discrimination ability of comorbidity, frailty, and subjective health to predict mortality in community-dwelling older people: Population based prospective cohort study. *Eur J Intern Med* 2017; **42**:29-38.
- 131 Lai FTT, Guthrie B, Wong SYS, et al. Sex-specific intergenerational trends in morbidity burden and multimorbidity status in hong kong community: An age-period-cohort analysis of repeated population surveys. *BMJ Open* 2019; **9**:1-10.
- 132 Lai FTT, Ma TW, Hou WK. Multimorbidity is associated with more subsequent depressive symptoms in three months: A prospective study of community-dwelling adults in hong kong. *Int Psychogeriatr* 2018; **31**:1367-71.
- 133 Lares PA Perelman J. The current and projected burden of multimorbidity: A cross-sectional study in a southern europe population. *Eur J Ageing* 2019; **16**:181-92.

- 134 Lang K, Alexander IM, Simon J, et al. The impact of multimorbidity on quality of life among midlife women: Findings from a u.S. Nationally representative survey. *J Women's Health* 2015; **24**:374-83.
- 135 Le Cossec C, Perrine AL, Beltzer N, Fuhrman C, Carcaillon-Bentata L. Pre-frailty, frailty, and multimorbidity: Prevalences and associated characteristics from two french national surveys. *J Nutr Health Aging* 2016; **20**:860-69.
- 136 Lee TA, Shields AE, Vogeli C, et al. Mortality rate in veterans with multiple chronic conditions. *J Gen Intern Med* 2007; **22** 403-7.
- 137 Lee WJ, Peng LN, Lin CH, Lin HP, Loh CH, Chen LK. The synergic effects of frailty on disability associated with urbanization, multimorbidity, and mental health: Implications for public health and medical care. *Sci Rep* 2018; **8**:1-7.
- 138 Li J, Green M, Kearns B, et al. Patterns of multimorbidity and their association with health outcomes within yorkshire, england: Baseline results from the yorkshire health study. *BMC Public Health* 2016; **16**:1-9.
- 139 Li Y, White K, O'Shields KR, McLain AC, Merchant AT. Light-intensity physical activity and cardiometabolic risk among older adults with multiple chronic conditions. *Am J Health Promot* 2019; **33**:507-15.
- 140 Lujic S, Simpson JM, Zwar N, Hosseinzadeh H, Jorm L. Multimorbidity in australia: Comparing estimates derived using administrative data sources and survey data. *PLoS One* 2017; **12**:1-18.
- 141 Lupiáñez-Villanueva F, Anastasiadou D, Codagnone C, Nuño-Solinís R, Garcia-Zapirain Soto MB. Electronic health use in the european union and the effect of multimorbidity: Cross-sectional survey. *J Med Internet Res* 2018; **20**:1-14.
- 142 Zhou C-h, Tang S-f, Wang X-h, et al. Satisfaction about patient-centeredness and healthcare system among patients with chronic multimorbidity. *Current Med Sci* 2018; **38**:184-90.
- 143 Zhang R, Lu Y, Shi L, Zhang S, Chang F. Prevalence and patterns of multimorbidity among the elderly in china: A cross-sectional study using national survey data. *BMJ Open* 2019; **9**:1-9.
- 144 Wong CH, Weiss D, Sourial N, et al. Frailty and its association with disability and comorbidity in a community-dwelling sample of seniors in montreal: A cross-sectional study. *Aging Clin Exp Res* 2010; **22**:54-62.
- 145 Weimann A, Dai D, Oni T. A cross-sectional and spatial analysis of the prevalence of multimorbidity and its association with socioeconomic disadvantage in south africa: A comparison between 2008 and 2012. *Soc Sci Med* 2016; **163**:144-56.
- 146 Wang L, Palmer AJ, Cocker F, Sanderson K. Multimorbidity and health-related quality of life (hrqol) in a nationally representative population sample: Implications of count versus cluster method for defining multimorbidity on hrqol. *Health Qual life Outcomes* 2017; **15**:1-12.
- 147 Wang C, Pu R, Li Z, et al. Subjective health and quality of life among elderly people living with chronic multimorbidity and difficulty in activities of daily living in rural South Africa. *Clin Interv Aging* 2019; **14**:1285-96.
- 148 Wade A, Hobbs M, Green MA. Investigating the relationship between multimorbidity and dental attendance: A cross-sectional study of uk adults. *British Dental Journal* 2019; **226**:138-43.
- 149 Maciejewski ML Hammill BG. Measuring the burden of multimorbidity among medicare beneficiaries via condition counts and cumulative duration. *Health Serv Res* 2019; **54**:484-91.
- 150 Marengoni A, Angleman S, Meinow B, et al. Coexisting chronic conditions in the older population: Variation by health indicators. *Eur J Intern Med* 2016; **31**:29-34.

- 151 Marengoni A, von Strauss E, Rizzuto D, Winblad B, Fratiglioni L. The impact of chronic multimorbidity and disability on functional decline and survival in elderly persons. A community-based, longitudinal study. *J Intern Med* 2009; **265**:288-95.
- 152 Marques A, Santos DA, Peralta M, Sardinha LB, González Valeiro M. Regular physical activity eliminates the harmful association of television watching with multimorbidity. A cross-sectional study from the european social survey. *Prev Med* 2018; **109**:28-33.
- 153 Mavaddat N, Valderas JM, van der Linde R, Khaw KT, Kinmonth AL. Association of self-rated health with multimorbidity, chronic disease and psychosocial factors in a large middle-aged and older cohort from general practice: A cross-sectional study. *BMC Fam Pract* 2014; **15**:1-11.
- 154 McDaid O, Hanly MJ, Richardson K, Kee F, Kenny RA, Savva GM. The effect of multiple chronic conditions on self-rated health, disability and quality of life among the older populations of northern ireland and the republic of ireland: A comparison of two nationally representative cross-sectional surveys. *BMJ Open* 2013; **3**:1-9.
- 155 Melis R, Marengoni A, Angleman S, Fratiglioni L. Incidence and predictors of multimorbidity in the elderly: A population-based longitudinal study. *PLoS One* 2014; **9**:1-8.
- 156 Min LC, Wenger NS, Fung C, et al. Multimorbidity is associated with better quality of care among vulnerable elders. *Med Care* 2007; **45**:480-8.
- 157 Momtaz YA, Hamid TA, Yahaya N, Ibrahim R. Effects of chronic comorbidity on psychological well-being among older persons in northern peninsular Malaysia. *Appl Res Qual Life* 2010; **5**:133-46.
- 158 Mondor L, Cohen D, Khan AI, Wodchis WP. Income inequalities in multimorbidity prevalence in ontario, canada: A decomposition analysis of linked survey and health administrative data. *Int J Equity Health* 2018; **17**:1-13.
- 159 Muggah E, Graves E, Bennett C, Manuel DG. The impact of multiple chronic diseases on ambulatory care use; a population based study in ontario, canada. *BMC Health Serv Res* 2012; **12**:1-6.
- 160 Nagel A, Jungert A, Spinneker A, Neuhäuser-Berthold M. The impact of multimorbidity on resting metabolic rate in community-dwelling women over a ten-year period: A cross-sectional and longitudinal study. *J Nutr Health Aging* 2017; **21**:781-86.
- 161 Niedzwiedz CL, Katikireddi SV, Pell JP, Smith DJ. Sex differences in the association between salivary telomere length and multimorbidity within the us health & retirement study. *Age Ageing* 2019; **48**:703-10.
- 162 Nunes BP, Camargo-Figuera FA, Guttier M, et al. Multimorbidity in adults from a southern brazilian city: Occurrence and patterns. *Int J Public Health* 2016; **61**:1013-20.
- 163 Nunes BP, Chiavegatto Filho ADP, Pati S, et al. Contextual and individual inequalities of multimorbidity in brazilian adults: A cross-sectional national-based study. *BMJ Open* 2017; **7**:1-12.
- 164 Nunes BP, Thumé E, Facchini LA. Multimorbidity in older adults: Magnitude and challenges for the brazilian health system. *BMC Public Health* 2015; **15**:1-11.
- 165 Olaya B, Domènech-Abella J, Moneta MV, et al. All-cause mortality and multimorbidity in older adults: The role of social support and loneliness. *Exp Gerontol* 2017; **99**:120-26.
- 166 Olivares DEV, Chambi FRV, Chañi EMM, Craig WJ, Pacheco SOS, Pacheco FJ. Risk factors for chronic diseases and multimorbidity in a primary care context of

- central argentina: A web-based interactive and cross-sectional study. *Int J Environ Res Public Health* 2017; **14**:1-22.
- 167 Park B, Ock M, Lee HA, et al. Multimorbidity and health-related quality of life in koreans aged 50 or older using knhanes 2013-2014. *Health Qual Life Outcomes* 2018; **16**:1-10.
- 168 Patel KV, Peek MK, Wong R, Markides KS. Comorbidity and disability in elderly mexican and mexican american adults: Findings from mexico and the southwestern united states. *J Aging Health* 2006; **18**:315-29.
- 169 Pati S, Hussain MA, Swain S, et al. Development and validation of a questionnaire to assess multimorbidity in primary care: An indian experience. *BioMed Res Int* 2016; **2016**:1-9.
- 170 Pati SS, Subhashisa: Knottnerus, J.: Metsemakers, Job F.: van den Akker, Marjan. Health related quality of life in multimorbidity: A primary-care based study from odisha, India. *Health Qual Life Outcomes* 2019; **17**:1-11.
- 171 Payne RA, Abel GA, Guthrie B, Mercer SW. The effect of physical multimorbidity, mental health conditions and socioeconomic deprivation on unplanned admissions to hospital: A retrospective cohort study. *Can Med Assoc J* 2013; **185**:E221-8.
- 172 Pérez LM, Hooshmand B, Mangialasche F, et al. Glutathione serum levels and rate of multimorbidity development in older adults. *J Gerontol A Biol Sci Med Sci* 2020; **75**:1089-94.
- 173 Petersen I, Rathod S, Kathree T, Selohilwe O, Bhana A. Risk correlates for physical-mental multimorbidities in south africa: A cross-sectional study. *Epidemiol Psychiatr Sci* 2019; **28**:418-26.
- 174 Pfortmueller CA, Stotz M, Lindner G, Müller T, Rodondi N, Exadaktylos AK. Multimorbidity in adult asylum seekers: A first overview. *PloS one* 2013; **8**:1-7.
- 175 Pressley JC Patrick CH. Frailty bias in comorbidity risk adjustments of community-dwelling elderly populations. *J Clin Epidemiol* 1999; **52**:753-60.
- 176 Prior A, Fenger-Grøn M, Larsen KK, et al. The association between perceived stress and mortality among people with multimorbidity: A prospective population-based cohort study. *Am J Epidemiol* 2016; **184**:199-210.
- 177 Ribeiro AS, Pereira LC, Silva DRP, et al. Physical activity and sitting time are specifically associated with multiple chronic diseases and medicine intake in brazilian older adults. *J Aging Phys Act* 2018; **26**:608-13.
- 178 Ruel G, Lévesque J-F, Stocks N, et al. Understanding the evolution of multimorbidity: Evidences from the north west adelaide health longitudinal study (nwaHS). *PLOS ONE* 2014; **9**:1-11.
- 179 Ruel G, Shi Z, Zhen S, et al. Association between nutrition and the evolution of multimorbidity: The importance of fruits and vegetables and whole grain products. *Clin Nutr* 2014; **33**:513-20.
- 180 Ryan A, Murphy C, Boland F, Galvin R, Smith SM. What is the impact of physical activity and physical function on the development of multimorbidity in older adults over time? A population-based cohort study. *J Gerontol A Biol Sci Med Sci* 2018; **73**:1538-44.
- 181 Schmidt AE, Ilinca S, Schulmann K, et al. Fit for caring: Factors associated with informal care provision by older caregivers with and without multimorbidity. *Eur J Ageing* 2016; **13**:103-13.
- 182 Schöttker B, Saum KU, Jansen EH, Holleczeck B, Brenner H. Associations of metabolic, inflammatory and oxidative stress markers with total morbidity and multimorbidity in a large cohort of older german adults. *Age Ageing* 2016; **45**:127-35.

- 183 Seo JC, B.: Kim, S.: Lee, H.: Oh, D. The relationship between multiple chronic diseases and depressive symptoms among middle-aged and elderly populations: Results of a 2009 Korean community health survey of 156,747 participants. *BMC Public Health* 2017; **17**:1-10.
- 184 She R, Yan Z, Jiang H, Vetrano DL, Lau JTF, Qiu C. Multimorbidity and health-related quality of life in old age: Role of functional dependence and depressive symptoms. *J Am Med Dir Assoc* 2019; **20**:1143-49.
- 185 Singh K, Patel SA, Biswas S, et al. Multimorbidity in South Asian adults: Prevalence, risk factors and mortality. *J Public Health (Oxf)* 2019; **41**:80-89.
- 186 Spangenberg L, Forkmann T, Brähler E, Glaesmer H. The association of depression and multimorbidity in the elderly: Implications for the assessment of depression. *Psychogeriatr* 2011; **11**:227-34.
- 187 Stickley A, Koyanagi A, Ueda M, Inoue Y, Waldman K, Oh H. Physical multimorbidity and suicidal behavior in the general population in the United States. *J Affect Disord* 2020; **260**:604-09.
- 188 Streit S, da Costa BR, Bauer DC, et al. Multimorbidity and quality of preventive care in Swiss university primary care cohorts. *PLOS ONE* 2014; **9**:1-7.
- 189 Stubbs B, Vancampfort D, Veronese N, et al. Multimorbidity and perceived stress: A population-based cross-sectional study among older adults across six low- and middle-income countries. *Maturitas* 2018; **107**:84-91.
- 190 Su P, Ding H, Zhang W, et al. The association of multimorbidity and disability in a community-based sample of elderly aged 80 or older in Shanghai, China. *BMC Geriatr* 2016; **16**:1-7.
- 191 Sundstrup E, Jakobsen MD, Mortensen OS, Andersen LL. Joint association of multimorbidity and work ability with risk of long-term sickness absence: A prospective cohort study with register follow-up. *Scand J Work Environ Health* 2017; **43**:146-54.
- 192 Takahashi PY, Ryu E, Hathcock MA, et al. A novel housing-based socioeconomic measure predicts hospitalisation and multiple chronic conditions in a community population. *J Epidemiol Community Health* 2016; **70**:286-91.
- 193 Tinetti MEM, G. J.: Chang, S. S.: Newman, A. B.: Fitzpatrick, A. L.: Fried, T. R.: Peduzzi, P. N. Contribution of multiple chronic conditions to universal health outcomes. *J Am Geriatr Soc* 2011; **59**:1686-91.
- 194 Troelstra SA, Straker L, Harris M, Brown S, van der Beek AJ, Coenen P. Multimorbidity is common among young workers and related to increased work absenteeism and presenteeism: Results from the population-based Raine Study cohort. *Scand J Work Environ Health* 2020; **46**:218-27.
- 195 van Zon SKR, Reijneveld SA, Galaurchi A, Mendes de Leon CF, Almansa J, Bültmann U. Multimorbidity and the transition out of full-time paid employment: A longitudinal analysis of the Health and Retirement Study. *J Gerontol B Psychol Sci Soc Sci* 2020; **75**:705-15.
- 196 Vancampfort D, Stubbs B, Koyanagi A. Physical chronic conditions, multimorbidity and sedentary behavior amongst middle-aged and older adults in six low- and middle-income countries. *Int J Behav Nutr Phys Act* 2017; **14**:1-13.
- 197 Vassilaki M, Aakre JA, Cha RH, et al. Multimorbidity and risk of mild cognitive impairment. *J Am Geriatr Soc* 2015; **63**:1783-90.
- 198 Vassilaki M, Aakre JA, Mielke MM, et al. Multimorbidity and neuroimaging biomarkers among cognitively normal persons. *Neurology* 2016; **86**:2077-84.
- 199 Villarreal AEG, S.: Lopez, L.: Britton, G. B.: Initiative, P. A. R. Cognitive impairment, depression, and co-occurrence of both among the elderly in Panama:

- Differential associations with multimorbidity and functional limitations. *BioMed Res Int* 2015; **2015**:1-7.
- 200 Violán C, Foguet-Boreu Q, Fernández-Bertolín S, et al. Soft clustering using real-world data for the identification of multimorbidity patterns in an elderly population: Cross-sectional study in a mediterranean population. *BMJ Open* 2019; **9**:1-14.
- 201 von Strauss E, Fratiglioni L, Viitanen M, Forsell Y, Winblad B. Morbidity and comorbidity in relation to functional status: A community-based study of the oldest old (90+ years). *J Am Geriatr Soc* 2000; **48**:1462-9.
- 202 Vos HM, Bor HH, Rangelrooij-Minkels MJ, Schellevis FG, Lagro-Janssen AL. Multimorbidity in older women: The negative impact of specific combinations of chronic conditions on self-rated health. *Eur J Gen Pract* 2013; **19**:117-22.
- 203 Vu HMN, L. H.: Tran, T. H.: Pham, K. T. H.: Phan, H. T.: Nguyen, H. N.: Tran, B. X.: Latkin, C. A.: Ho, C. S. H.: Ho, R. C. M. Effects of chronic comorbidities on the health-related quality of life among older patients after falls in vietnamese hospitals. *Int J Environ Res Public Health* 2019; **16**:1-9.
- 204 Wang L, Cocker F, Kilpatrick M, et al. The associations of multimorbidity with health-related productivity loss in a large and diverse public sector setting: A cross-sectional survey. *J Occup Environ Med* 2018; **60**:528-35.
- 205 Wang X-X, Lin W-Q, Chen X-J, et al. Multimorbidity associated with functional independence among community-dwelling older people: A cross-sectional study in southern china. *Health Qual Life Outcomes* 2017; **15**:73-9.
- 206 Wijers IGM, Ayala A, Rodriguez-Blazquez C, et al. The disease burden morbidity assessment in older adults and its association with mortality and other health outcomes. *Eur J Ageing* 2019; **16**:193-203.
- 207 Williams JSE, L. E. The association between multimorbidity and quality of life, health status and functional disability. *Am J Med Sci* 2016; **352**:45-52.
- 208 Woldeamayyat EM, Kassa A, Gari T, Dangisso MH. Chronic diseases multimorbidity among adult patients at hawassa university comprehensive specialized hospital. *BMC Public Health* 2018; **18**:1-8.
- 209 Yao SS, Cao GY, Han L, et al. Associations between somatic multimorbidity patterns and depression in a longitudinal cohort of middle-aged and older chinese. *J Am Med Dir Assoc* 2020; **21**:1282-87.
- 210 Yorke AM, Curtis AB, Shoemaker M, Vangsnes E. The impact of multimorbidity on grip strength in adults age 50 and older: Data from the health and retirement survey (hrs). *Arch Gerontol Geriatr* 2017; **72**:164-68.
- 211 You L, Yu Z, Zhang X, et al. Association between multimorbidity and depressive symptom among community-dwelling elders in eastern china. *Clin Interv Aging* 2019; **14**:2273-80.
- 212 Zhang L, Ma L, Sun F, Tang Z, Chan P. A multicenter study of multimorbidity in older adult inpatients in china. *J Nutr Health Aging* 2020; **24**:269-76.
- 213 Khanam MA, Streatfield PK, Kabir ZN, Qiu C, Cornelius C, Wahlin Å. Prevalence and patterns of multimorbidity among elderly people in rural bangladesh: A cross-sectional study. *J Health, Popul Nutr* 2011; **29**:406-14.
- 214 Cornell J, Pugh J, Williams, Jr., et al. Multimorbidity clusters: Clustering binary data from multimorbidity clusters: Clustering binary data from a large administrative medical database. *Appl Multivar Res* 2009; **12**:1-6.
- 215 Cassell A, Edwards D, Harshfield A, et al. The epidemiology of multimorbidity in primary care: A retrospective cohort study. *Br J GenPract* 2018; **68**:e245-e51.

- 216 Wong E Ly, Xu RH, Cheung AWI. Measuring the impact of chronic conditions and associated multimorbidity on health-related quality of life in the general population in hong kong sar, china: A cross-sectional study. *PLOS ONE* 2019; **14**:1-16.
- 217 Puth M-T, Weckbecker K, Schmid M, Münster E. Prevalence of multimorbidity in germany: Impact of age and educational level in a cross-sectional study on 19,294 adults. *BMC Public Health* 2017; **17**:1-7.